



Diversity and Abundance of Avian Fauna in Cultivated and Riverine Habitats of Bahawalnagar, Pakistan

Javed Iqbal¹, Sajid Hussain^{*1}, Huma Azam¹, Nimra¹, Shanza Karamat¹, Hafiz Toheed¹ and Fareeha Nawaz¹

¹Department of Zoology, GC University Faisalabad, Punjab, Pakistan

*Correspondence: sajid612gb@gmail.com

ARTICLE INFO

ARTICLE HISTORY: CVJ-25-428

Received: 06 February 2025

Revised: 13 March 2025

Accepted: 26 April 2025

Published online: 29 April 2025

Key words:

Avian fauna
Bahawalnagar
Cultivated land
Diversity
River

ABSTRACT

The avian fauna of Punjab, Pakistan's District Bahawalnagar was examined between October 2022 and May 2023. District Bahawalnagar, which lies close to the Sutlej River, provides an excellent habitat for a variety of bird species. This study is the first survey of bird species carried out in Bahawalnagar. The landscapes of Bahawalnagar were separated into two groups: the riverine area and the agricultural region. Potential bird habitats were found in both the riverine and cultivated sectors. The birds were viewed using a binocular telescope with the naked eye. For odd time periods, the survey was carried out using the point count method from a fixed location inside a specified 50-meter radius. 10515 individual birds of 74 species, spanning 32 families of 12 orders, were identified during the current study. Additionally, 25 plant species were found in the study region. Winter observations included 49 bird species, spring observations included 70, and summer observations included 60. Common Myna (*Acridotheres tristis*), House Crow (*Corvus splendens*) and House Sparrow (*Passer domesticus*) were the most prevalent species. According to statistical analysis, the spring season's evenness and Shannon-Wiener values were 0.6477 and 1.195, respectively. Species diversity was much higher in the spring than in other seasons. The order Passeriformes was the most prevalent in all food kinds and species diversity. These results indicate that agricultural and riverine habitats support diverse avian fauna. These approaches help support habitat diversity and food availability.

To Cite This Article: Iqbal J, Hussain S, Azam H, Nimra, Karamat S, Toheed H and Nawaz F, xxxx. Diversity and abundance of avian fauna in cultivated and riverine habitats of Bahawalnagar, Pakistan. Continental Vet J, x(x): xxx <http://dx.doi.org/10.71081/cvj/2025.038>

INTRODUCTION

Birds are the commonest species of vertebrates over the world with almost 10,400 species and making a great indicator of ecosystems (Barrowclough et al. 2016). Pakistan provides a platform of great climatic and vegetation regions that in means a large number of bird's species. To prevent from a cold winter season, large number of birds is migrating from Europe and Central Asia to Pakistan in winter (Stephens et al. 2016). Along with a bird, man-made landscape modifications typically result in a decrease in the natural environment and a subsequent loss of biological diversity. Birds are an important part of the global biodiversity and are essential to the upkeep of a healthy ecosystem. With their behavior, exquisite colors, and melodies, birds are thought to be human. Riparian regions also consist of a dispersal pathway and provide the enough cover for migratory birds; hence they also support a richness species of bird's diversity (Fraixedas et al. 2020). Vultures and the pied

crow (*Corvus albus*) are venomous birds that consume and decompose disease-carrying insects. In addition to controlling illness, birds are crucial for pollination and seed dissemination. Bird diversity and abundance are greatly enhanced by agro ecosystems. Small rodents that permanently harm crops are managed by birds (Chain-Guadarrama et al. 2019).

The world's fifth most populous nation is Pakistan. Bird diversity is thus seriously threatened by overpopulation brought on by habitat loss, deforestation, altered desert ecosystems, industrialization, urbanization, and agricultural practices. Birds that rely on the forest have been primarily harmed by its degradation, and when a forest fragment is too small or too degraded to guide prospective populations, reasons of species extinction have been hypothesized (Umar et al. 2018). Determining avian diversity through regional studies is crucial to species protection. Birds are excellent environmental indicators that aid in the analysis of high-quality conservation sites. Pakistan's birds are seriously

The use of pesticides, fertilizers, shorter crop rotations, machine-driven farming, and making agricultural fields less suitable for a wide variety of organisms all contribute to the intensification that takes place at the field level (Moreau et al. 2022). Numerous ecosystems can be found in the region, including the river, woodlands, desert, and both farmed and uncultivated land. The wild plant *Tamarix dioica*, also referred to as tamarisk, which formerly covered a sizable amount of the

MATERIALS AND METHODS

The Sutlej River runs along the northern boundary of the Bahawalnagar district (29.6892° N, 72.9933° E), which is bounded to the east and south by Indian Territory and to the west by the Bahawalpur district. It is situated 163 meters (516.41 feet) above sea level and is located on the expanding flat plains of Southern Punjab. District Bahawalnagar has a total size of 8,878 km². It is the 54th largest city in Pakistan by population, according to the 2017 census. According to Pakistan's 2017 census, Bahawalnagar had 2,981,919 residents. Bahawalnagar supports a range of habitats. Two studies sites—Site 1 in the riverine region and Site 2 in the cultivated area—were built in the riverine and cultivated areas, which served as the two primary research areas (Fig. 1).




Fig. 1: Map of District Bahawalnagar indicating the study sites of riverine and agriculture sites field Survey.

Field Surveys

The period of the field observation was October 2022–May 2023. Three distinct seasons—winter, spring, and summer—were used to gather data. The point count method was used to gather data from a certain location for a predetermined amount of time. This approach was effective for studying the species in a wide variety of environments. The point count sampling approach was used to record birds at several points within a predetermined radius (Morelli et al. 2022). Five plots for each location in each season were placed 300 meters apart to create 105-point count stations spread across six habitats. A global positioning system (GPS) device was used to create plots with a 50-meter radius. For every kind of micro landscape, five plots were randomly chosen inside a five-kilometer-by-five-kilometer grid cell (Gasc et al. 2015). Every plot was observed for twenty minutes. Every plot was visited twice during the day, with equal intervals of two hours at daybreak and dusk. The bird diversity (total/physical count) along the river was determined using the linear count survey method (0.4 km in length and 0.2 km in breadth). The birds were photographed with a camera (Canon EOS 5D), binoculars, and naked eyes. A number of helpful bird species identification aids were examined (Sullivan et al. 2017).

Statistical Analysis

To analyze avian diversity across different habitats or districts, the following ecological indices were applied:

1. Relative Abundance (Pi)

Relative abundance was used to determine the proportion of individual bird species relative to the total bird population in each site:

$$\text{Relative Abundance (Pi)} = N/n$$

Where:

- n = Number of individuals of a specific bird species in a site
- N = Total number of individuals of all bird species in that site

2. Shannon-Wiener Diversity Index (H')

This index was used to quantify species diversity by considering both richness and evenness of the species present:

$$H' = -\sum (P_i \cdot \ln P_i)$$

Where:

- H' = Shannon-Wiener diversity index
- P_i = Proportion of each species relative to the total number of individuals
- $\ln P_i$ = Natural logarithm of P_i

3. Species Richness (S) and Evenness Index (J')

Species richness refers to the total number of different bird species recorded in a given area:

$$S = \text{Total number of species}$$

Evenness (J') was calculated to assess how evenly individuals are distributed among the species:

$$J' = H'/H_{\max} = H'/\ln S$$

Where:

- J' = Evenness index
- H' = Shannon-Wiener index

- H_{\max} = Maximum diversity possible = $\ln(S)$

4. Simpson's Diversity Index (D)

This index was used to measure the probability that two individuals randomly selected from a sample belong to different species:

$$D = N-1 \sum (n-1)/N - 1$$

Where:

- n = Number of individuals of a specific bird species
- N = Total number of individuals of all species

Vegetation Assessment

At bird counting stations, the quadrant technique (10m 10m) was used to sample the plant variables present in these ecosystems. This approach is one of the most popular and well-respected ways to survey vegetation in different settings (Besnard et al. 2015). To ascertain the vegetation structure within the consistency of the point count stations, a total of 21 quadrant plots were gathered. Plant pressure was assessed using field-collected plant samples, according to (Agra et al. 2015). The dried plant samples were prepared, oven-dried for a week, and then stored on herbarium sheets. Plants from the Botany Department of Government College Faisalabad were identified.

RESULTS

In Bahawalnagar, 74 bird species from 32 families and 12 orders have been identified throughout the study period. There were 32 species and 13 families in the main order, Passeriformes. Forty different plant species were observed. Common mulberries (*Morus alba*), shisham (*Dalbergia sisso*), eucalyptus trees (*Eucalyptus globulus*), Bermuda grass (*Cynodon dactylon*), neem trees (*Azadirachta indica*), populus/cottonwood (*Populus tremula*), and sukhchain/Indian beach trees (*Pongamia pinnata*) were abundant. These plants were used by birds for roosting, nesting, and feeding. Certain plants offer bird's additional safety. Over the course of the study, there was a notable diversity of bird species in the Bahawalnagar area (70 spring, 25 winter, and 25 summer species) Fig 3. The avian diversity index for the area was 2.25. Along with their own unique species, each of the two research sites also featured a few common species that are found all over the world. Except for five species, every species observed during the investigation is of the least concern. IUCN Red Lists indicate that these five species are threatened (Table 1).

Table 1: IUCN conservation status of avian species recorded in Bahawalnagar

Sr.#	Species name	IUCN Status	Residential status
1	Spotted green shank	Endangered	E
2	Sandpiper	Critical	CR
3	Red Wattled Lapwing	Endangered Vulnerable	VU
4	Piping Plover	Near Threatened	NT
5	Pied Kingfisher	Vulnerable	VU

Study Site 1: The riverine area

Study Site 1 covered around 20 km² of the Satluj River. Along the riverbanks here are native plants including River Tamarind (*Leucaena leucocephala*), Common Mulberry (*Morus alba*), Common Wheat (*Triticum aestivum*), Berseem Clover (*Trifolium alexandrinum*), Shisham (*Dalbergia sissoo*), and Wild Sugarcane (*Saccharum spontaneum*). The river also features several sandy islands throughout its course. This location is home to a broad range of bird species throughout the year, with the maximum avian richness occurring in the winter. A total of 5,606 individual birds, representing 53 distinct species, were recorded throughout the surveys, demonstrating the region's significant species diversity. February and March of 2022 had the greatest number of persons. The findings showed that the recorded avifauna belonged to nine orders and 26 families (Fig. 1). With 11 families and 19 species, the order Passeriformes was the most prevalent among them. With five families and fifteen species, the Charadriiformes order was the second most represented. The order Coraciiformes is the third most represented, with three families and four species. The most common species at Study Site 1 was the House Crow (*Corvus splendens*), which belongs to the Passeriformes order and had 1,213 individuals documented, or a relative abundance (RA) of 5.35%.

The Great Blue Heron (*Ardea herodias*), a member of the order Pelecaniformes, was the least frequent species, with only two individuals observed (RA 0.0088%). The Black-winged Stilt (*Himantopus himantopus*, RA 1.9722%), Common Myna (*Acridotheres tristis*, RA 2.2855%), and House Crow (RA 5.352%) were the most common species at Study Site 2. Other types of terrain also showed seasonal variations in bird abundance. Seasonal variations in bird counts were seen in the riverine zones, with 953 individuals recorded in the winter, 3,241 in the spring, and 1,022 in the summer. The study site has a high degree of bird diversity, according to the diversity measures. Table 2 shows that the Simpson Diversity Index (D) was 0.0865, the Evenness (J') value was 0.7407, and the Shannon-Wiener Diversity Index (H') was 1.2772. The first-time documentation of three bird species in District Bahawalnagar—the Glossy Ibis (*Plegadis falcinellus*), the Short-tailed Pipit (*Anthus brachyurus*), and the Red-naped Ibis (*Pseudibis papillosa*)—was a noteworthy highlight of this study. These species, which are mostly known to exist in Indian Territory and the Pakistani province of Sindh, have never before been documented in the area. Their known distribution range has been significantly expanded by their presence in Bahawalnagar.

Study Site 2: The cultivated area

In addition to orange and guava orchards, the area is home to a wide range of agricultural crops, such as vegetables, maize, rice, wheat, and sugarcane. At this

location, 4,909 individual birds were counted, representing 45 species richness. Ten orders and twenty-six families were used to categorize the avifaunal data. With 13 families and 23 species, the Passeriformes order was the most prevalent. The order Coraciiformes, which had three families and three species, came next. With two families and two species apiece, the orders Charadriiformes and Bucerotiformes were tied for third place. The most common species at this site was the House Crow (*Corvus splendens*), a species of the Passeriformes order, with 1,273 individuals recorded. The Black-crowned Night Heron (*Nycticorax nycticorax*), a species of the Pelecaniformes order, was the least common, with only one bird observed. In 2023, the highest number of persons was in February and March. There was a noticeable seasonal variation in bird abundance, with 861 birds recorded in the winter, 2,311 in the spring, and 972 in the summer. House Crow (RA 5.617%), Common Myna (*Acridotheres tristis*, RA 4.4917%), and House Sparrow (*Passer domesticus*, RA 4.0019%) were the most prevalent species at Site 2 (Table 2). With a Simpson Diversity Index (D) of 0.1529, an Evenness (J') of 0.5884, and a Shannon-Wiener Index (H') of 0.9728, the diversity indices showed a reasonably high degree of avian variety.

3. The entire study area:

Bird surveys covering three different seasons—winter (October, November, December, and January), spring (February and March), and summer (April and up to May 18)—were carried out between October 2022 and May 2023 in order to evaluate seasonal variance. The survey period was two months for the spring and summer seasons, and four months for the winter season. To provide thorough coverage, every study location was visited at random during each season (Fig. 2). Seasonal temperature variations were evident in the climate data collected during the research period. The lowest average temperature in June was 27°C (82°F), while the highest average temperature was 41°C (105°F). On the other hand, January's average temperature was as low as 4°C (39°F) and as high as 20°C (68°F). In all seasons, 74 different bird species were observed during the study period. Over the winter, 49 species were seen, with an Evenness (J') of 0.6961 and a Shannon-Wiener Diversity Index (H') of 1.1766. The corresponding diversity values were H' = 1.195 and J' = 0.6477 when the number of species observed increased to 70 in the spring. Of the 60 species discovered during the summer, the highest diversity was observed with H' = 1.3022 and J' = 0.7323 (Table 3). Across all survey sites, the most common and frequently sighted species were the House Sparrow (*Passer domesticus*), Black-winged Stilt (*Himantopus himantopus*), Common Myna (*Acridotheres tristis*), and House Crow (*Corvus splendens*).

Table 2: Checklist of birds recorded in Agriculture area and around Sutlej River

Order	Family	Scientific Name	Common Name	Site 1	Site 2	RA
Passeriformes	Covidae	<i>Corvus splendens</i>	House Crow	1213	1273	10.969
		<i>Pica hudsonia</i>	Black billed magpie	0	14	0.0618
		<i>Acridotheres tristis</i>	Common Myna	518	1018	6.7772
	Pycnonotidae	<i>Pycnonotus cafer</i>	Red vented bulbul	3	201	0.9
	Laniidae	<i>Lanius vittatus</i>	Bay backed shrike	0	53	0.2338

		<i>Lanues schach</i>	Long tailed shrike	0	13	0.0026
	Passeridae	<i>Passer domesticus</i>	House Sparrow	102	907	4.4519
	Motacillidae	<i>Anthus brachyurus</i>	Short tailed Pipit	0	7	0.0014
		<i>Anthus rubescens</i>	Buff bellied pipit	29	0	0.1279
		<i>Motacilla flava</i>	Western yellow wagtail	102	11	0.4985
		<i>Anthus similis</i>	Long billed Pipit	6	0	0.0265
		<i>Motacilla citreola</i>	Citrine Wagtail	109	16	0.5515
		<i>Motacilla maderaspatensis</i>	White browed wagtail	17	0	0.0750
		<i>Anthus Novaeseelandiae</i>	Australasian Pipit	6	0	0.0265
		<i>Motacilla alba</i>	White wagtail	387	7	1.7075
	Leiothrichidae	<i>Argya striata</i>	Jungle babbler	0	61	0.2691
	Muscicapidae	<i>Copsychus saularis</i>	Oriental Magpie robin	291	7	1.3149
		<i>Oenanthe fusca</i>	Brown rock Chat	11	42	0.2338
	Passerellidae	<i>Melospiza fusca</i>	Canyon towhee	0	33	0.1456
		<i>Peucaea cassinii</i>	Cassins Sparrow	9	0	0.0397
	Hirundinidae	<i>Riparia riparia</i>	Bank swallow	123	0	0.5427
	Cisticolidae	<i>Prinia gracilis</i>	Graceful prinia	6	12	0.0265
		<i>Prinia buehneri</i>	Rufous fronted prinia	0	31	0.1368
		<i>Prinia subflava</i>	Tawny flanked Prinia	0	11	0.0485
	Alaudidae	<i>Alauda arvensis</i>	Eurasian skylark	0	13	0.0573
		<i>Spizocorys conirostris</i>	Pink billed lark	0	19	0.0838
		<i>Eremophila alpestris</i>	Shore lark	13	0	0.0573
		<i>Galerida cristata</i>	Crested lark	39	41	0.353
	Acrocephalidae	<i>Iduna rama</i>	Sykes's warbler	0	11	0.0485
	Dicruridae	<i>Dicrurus macrocercus</i>	Black drongo	31	313	1.5178
Charadriiformes	Scolopacidae	<i>Tringa ochropus</i>	Green Sandpiper	50	0	0.2206
		<i>Calidris pusilla</i>	wader	40	0	0.1765
		<i>Numenius arquata</i>	Common curlew	11	0	0.0485
		<i>Tringa guttifer</i>	Spotted greenshank	19	0	0.0838
		<i>Tringa glareola</i>	Wood Sandpiper	9	0	0.0397
		<i>Tringa semipalmata</i>	Willet	9	0	0.0397
		<i>Calidris alpina</i>	Sandpiper	53	0	0.2338
		<i>Tringa nebularia</i>	Common greenshank	441	0	1.9458
	Charadriidae	<i>Charadrius vociferus</i>	Killdeer	54	0	0.2383
		<i>Vanellus indicus</i>	Red Wattled Lapwing	285	123	1.8002
		<i>Charadrius melodus</i>	Piping Plover	33	0	0.1456
		<i>Charadrius melodus</i>	Piping Plover	58	0	0.2559
	Recurvirostridae	<i>Himantopus himantopus</i>	Black Winged Stilt	447	12	2.0251
	Glareolidae	<i>Glareola pratincola</i>	Collared pratincole	76	0	0.3353
	Laridae	<i>Sterna Hirundo</i>	Common tern	357	0	1.5751
Galliformes	Phasianidae	<i>Coturnix coturnix</i>	Common Quail	24	13	0.1632
Coraciiformes	Alcedinidae	<i>Ceryle rudis</i>	Pied Kingfisher	120	0	0.5295
		<i>Halcyon smyrnensis</i>	White throated kingfisher	12	14	0.1147
	Coraciidae	<i>Coracias benghalensis</i>	Indian Roller	3	54	0.2515
		<i>Coracias benghalensis</i>	Indian Roller	29	24	0.2338
	Meropidae	<i>Merops persicus</i>	Blue cheeked bee eater	6	70	0.3353
Pelecaniformes	Threskiornithidae	<i>Pseudibis papillosa</i>	Red naped ibis	14	0	0.0618
		<i>Plegadis falcinellus</i>	Glossy ibis	11	0	0.0485
	Ardeidae	<i>Nycticorax nycticorax</i>	Black crowned night heron	7	1	0.0353
		<i>Ardea intermedia</i>	Intermediate Egret	16	0	0.0706
		<i>Bubulcus ibis</i>	Cattle egret	44	105	0.6574
		<i>Ardea herodias</i>	Great blue heron	2	0	0.0088
		<i>Egretta garzetta</i>	Little Egret	104	7	0.4898
		<i>Ardea alba</i>	Great Egret	210	139	1.5398
Anseriformes		<i>Anser indicus</i>	Bar headed Goose	11	0	0.0485
Cuculiformes	Cuculidae	<i>Eudynamis scolopacea</i>	Asian koel	0	17	0.0750
		<i>Centropus sinensis</i>	Greater coucal	0	29	0.1279
Columbiformes	Columbidae	<i>Streptopelia decaocta</i>	Eurasian collared dove	0	38	0.1677
		<i>Streptopelia tranquebarica</i>	Red collared Dove	16	25	0.1809
		<i>Columba livia domestica</i>	Feral pigeon	14	0	0.0618
		<i>Columba livia</i>	Rock pigeon	0	22	0.0970
		<i>Spilopelia chinensis</i>	columbinae	0	15	0.0662
		<i>Columbina livia domestica</i>	columbinae	0	21	0.0926
		<i>Spilopelia senegalensis</i>	Laughing dove	0	27	0.1191
Strigiformes	Strigidae	<i>Athene noctua</i>	Little owl	0	3	0.0132
Bucerotiformes	Bucerotidae	<i>Ocyrceros birostris</i>	Indian grey horn bill	0	23	0.1015
	Upupidae	<i>Upupa epops</i>	Hoopoe	3	6	0.0397
Piciformes	Picidae	<i>Dinopium benghalense</i>	Black rumped flame-back	0	7	0.0309
Gruiformes	Rallidae	<i>Amaurornis phoenicurus</i>	White breasted waterhen	3	0	0.0132
Total				5606	4909	46.2255




Fig. 2: Different orders of avian fauna demonstrating number of species.




Fig. 3: Seasonal distribution of bird fauna.




Fig. 4: Relative abundance of bird fauna of different sites.

Table 3: Diversity of avian fauna in Bahawalnagar

Diversity measure	Riverine area	Cultivated area	Total study area
Species richness	53	45	74
Shannon -Wiener (H')	1.2772	0.9728	2.25
Simpson diversity (D)	0.0865	0.1529	0.2394
Evenness (J')	0.7407	0.0865	1.3291

DISCUSSION

The diversity of birds in Punjab, Pakistan's District Bahawalnagar is described in detail for the first time in this study. This region had not previously been the focus of ornithological surveys, despite the ecological importance of the Sutlej River. The separation of the region into agricultural and riverine landscapes allowed for a methodical evaluation of the habitat-specific bird diversity. The vast number of species (74) and individuals (10,515) recorded indicates the availability and suitability of these habitats for supporting a diverse bird community. The order Passeriformes was shown to be the most dominating in all seasons, which is consistent with other studies that highlight the order's wide adaptability and diversity in a range of habitats. Due to their synanthropic tendencies and ability to thrive in ecosystems that people have altered, the most common species were House Crow, Common Myna, and House Sparrow. There was a clear seasonal fluctuation in species richness and diversity, with the maximum diversity indices occurring in the spring (Shannon-Wiener $H' = 1.195$; Evenness $J' = 0.6477$).

According to comparable observations in agro-riverine environments, this may be explained by migratory influx, reproductive activity, and an abundance of food resources during this season. The point count method's usefulness in avifaunal research was further supported by the fact that it was successful in sampling a range of habitats and producing accurate population estimates when applied within a defined 50-meter radius. Additionally, the discovery of uncommon species that had not been previously documented in this district, like *Plegadis falcinellus*, *Anthus brachyurus*, and *Pseudibis papillosa*, emphasizes the significance of ongoing observation and the promise of Bahawalnagar as a major location for bird conservation. However, the study also emphasizes the dangers that humans pose to the local avifauna, namely habitat loss, pollution, and hunting. These findings align with global concerns regarding the loss of biodiversity in riparian and agricultural habitats. Conservation efforts should focus on habitat preservation, increasing public awareness, and enforcing regulations in order to preserve the biological integrity of this region.

Conclusion

The study recorded 10,515 individual birds from 74 species, representing 32 families and 12 orders, along with 25 plant species. Bird observations were conducted both with binocular telescopes and the naked eye, using point count methods during odd time periods. Spring showed the highest species diversity and evenness, with Shannon-Wiener and evenness values of 1.195 and 0.6477, respectively. The most commonly observed birds were Common Myna, House Crow, and House Sparrow. Passeriformes emerged as the dominant order across all

types of habitats and seasons. The results indicate that cultivated land and riverine habitats diverse and species-rich bird community. These findings suggest that agricultural landscapes and riverine habitats can sustain diverse bird fauna.

REFERENCES

- Abrahamczyk, S 2019. Comparison of the ecology and evolution of plants with a generalist bird pollination system between continents and islands worldwide. *Biological Reviews*, 94(5): 1658-1671.
- Aebischer, N 2019. Fifty-year trends in UK hunting bags of birds and mammals, and calibrated estimation of national bag size, using GWCT's National Gamebag Census. *European Journal of Wildlife Research*, 65(4): 64.
- Agra, M, Bilenca, D, and Codesido, M 2015. Responses of birds to planting of *Lotus tenuis* pasture in the Flooding Pampas, Argentina. *Emu-Austral Ornithology*, 115(3): 270-276.
- Barrowclough, GF, Cracraft, J, Klicka, J, and Zink, RM 2016. How many kinds of birds are there and why does it matter? *PLoS One*, 11(11): e0166307.
- Basto, MN, Nicastro, KR, Tavares, AI, McQuaid, CD, Casero, M, Azevedo, F, and Zardi, GI 2019. Plastic ingestion in aquatic birds in Portugal. *Marine Pollution Bulletin*, 138: 19-24.
- Besnard, AG, Davranche, A, Maugenest, S, Bouzillé, J-B, Vian, A, and Secondi, J 2015. Vegetation maps based on remote sensing are informative predictors of habitat selection of grassland birds across a wetness gradient. *Ecological Indicators*, 58: 47-54.
- Carrasco L, Norton L, Henrys P, Siriwardena GM, Rhodes, CJ, Rowland C, and Morton, D, 2018. Habitat diversity and structure regulate British bird richness: Implications of non-linear relationships for conservation. *Biological Conservation*, 226: 256-263.
- Chain-Guadarrama, A, Martínez-Salinas, A, Aristizábal, N, and Ricketts, T H 2019. Ecosystem services by birds and bees to coffee in a changing climate: A review of coffee berry borer control and pollination. *Agriculture, Ecosystems and Environment*, 280: 53-67.
- de Montaigne CT, and Goulson D, 2023. Habitat quality, urbanisation and pesticides influence bird abundance and richness in gardens. *Science of the Total Environment*, 870: 161916.
- Fraixedas S, Lindén A, Piha M, Cabeza M, Gregory R, and Lehikoinen A, 2020. A state-of-the-art review on birds as indicators of biodiversity: Advances, challenges, and future directions. *Ecological Indicators*, 118: 106728.
- Gasc A, Pavoine S, Lellouch L, Grandcolas P and Sueur J, 2015. Acoustic indices for biodiversity assessments: Analyses of bias based on simulated bird assemblages and recommendations for field surveys. *Biological Conservation*, 191: 306-312.
- Iqbal F, Ayub Q, Wilson R, Song BK, Talei A, Yeong KY, Hermawan AA, Fahim M and Rahman S, 2021. Monitoring of heavy metal pollution in urban and rural environments across Pakistan using House crows (*Corvus splendens*) as bioindicator. *Environmental Monitoring and Assessment*, 193: 1-16.
- Matuoka MA, Benchimol M, de Almeida-Rocha JM and Morante-Filho JC, 2020. Effects of anthropogenic disturbances on bird functional diversity: A global meta-analysis. *Ecological Indicators*, 116: 106471.
- Moreau J, Rabdeau J, Badenhausser I, Giraudeau M, Sepp T, Crépin M, Gaffard A, Bretagnolle V and Monceau K, 2022. Pesticide impacts on avian species with special reference to farmland birds: a review. *Environmental Monitoring and Assessment*, 194(11): 790.
- Morelli F, Brlík V, Benedetti Y, Bussièrè R, Moudrá L, Reif J and Svitok M, 2022. Detection rate of bird species and what it depends on: Tips for field surveys. *Frontiers in Ecology and Evolution*, 9: 671492.
- Quintero I and Jetz W, 2018. Global elevational diversity and diversification of birds. *Nature*, 555(7695): 246-250.
- Saleem M, Javid A, Hussain A and Mehmood S, 2025. Diversity and molecular identification of bird species from South Punjab, Pakistan. *Journal of Wildlife and Biodiversity*, 9(1): 280-299.
- Stephens PA, Mason LR, Green RE, Gregory RD, Sauer JR, Alison J, Aunins A, Brotons L, Butchart SH and Campedelli T, 2016. Consistent response of bird populations to climate change on two continents. *Science*, 352(6281): 84-87.
- Sullivan BL, Phillips T, Dayer AA, Wood CL, Farnsworth A, Iliff MJ, Davies IJ, Wiggins A, Fink D and Hochachka WM, 2017. Using open access observational data for conservation action: A case study for birds. *Biological Conservation*, 208: 5-14.
- Umar M, Hussain M, Murtaza G, Shaheen FA and Zafar F, 2018. Ecological concerns of migratory birds in Pakistan: A review. *Punjab University Journal of Zoology*, 33(1): 69-76.
- Van Doren BM and Horton KG, 2018. A continental system for forecasting bird migration. *Science*, 361(6407): 1115-1118.