



Antimicrobial Potential of *Juglans regia* L (Walnut) Against Oral Pathogens Associated with Caries and Periodontitis

Azhar Rafique^{1*}, Muhammad Sharjeel Arshad², Nadeem Ali³ and Md. F. Kulyar⁴

¹Department of Zoology, Faculty of Life Sciences, Government College University Faisalabad; ²Institute of Microbiology, University of Agriculture, Faisalabad, Pakistan; ³Center of Excellence in Environmental Studies, King Abdulaziz University, Jeddah, Saudi Arabia; ⁴Department of Regenerative Medicine, State Research Institute Centre for Innovative Medicine, LT-08406 Vilnius, Lithuania

*Correspondence: azharrafique@gcuf.edu.pk

ARTICLE INFO

ARTICLE HISTORY: CVJ-26-209

Received: 15 January 2026
Revised: 20 February 2026
Accepted: 23 February 2026
Published online: 27 February 2026

Key words:

Walnut
Phenolic acids
Oral pathogens
Antimicrobial effects
Cytotoxicity

ABSTRACT

The non-transmissible diseases dental caries and ulatrophia pose a huge burden on human beings. Exploration for economical, easily accessible, harmless and substitute therapeutics development for the oral and dental health care is required. Natural products are used as traditional medicines from ancient times in Ayurvedic system of medicines. (walnut) is a type of plant that is commonly present around the Globe. The current research explores the antibacterial potential of three different solvent extracts including aqueous, acetone and methanol contrary to *Prevotella intermedia*, *Streptococcus mitis* and *Streptococcus mutans*. Vero cell line was used to determine the cellular cytotoxicity (CC₅₀). For the phytochemical analysis all three extracts were evaluated through HPLC (High Performance Liquid Chromatography). The findings of current experiment showed that the aqueous extract of have considerably higher antibacterial activity by measuring the diameter of zone of inhibition (20.00±0.58mm, 17.00±0.58mm) trailed by acetone and aqueous extracts counter to *S. mutans*, *S. mitis*, respectively. The methanolic extract proved statistically significant antibacterial activity at 160 mg/ml with inhibition zone (15.67±0.33mm) followed by acetone and aqueous extract against *P. intermedia*. The MIC (Minimum Inhibitory Concentration) (20mg/ml of aqueous extract against *S. mutans*; 5mg/ml of aqueous extract *S. mitis* and 10 mg/ml of methanol against *P. intermedia*). Highest quantity of phenolic acids was showed by methanol extract through HPLC while aqueous extract has considerably higher CC₅₀ and for therapeutics it's the safest among methanolic and acetone extract.

To Cite This Article: Rafique A, Arshad MS, Ali N and Kulyar MF, 2026. Antimicrobial potential of *Juglans regia* L (Walnut) against oral pathogens associated with caries and periodontitis. continental Vet J, 6(1): 118-125.
<http://dx.doi.org/10.71081/cvj/2026.070>

INTRODUCTION

Oral health is integral and vital for the human health, prosperity and well-being of life. Furthermore, the general health of an individual is interlinked with oral health; non-transmissible chronic diseases and oral cavity infections share “communal risk aspects” (Conway et al. 2013). Oral illnesses are most common factor that affecting the oral health, mostly inflammation of teeth (periodontitis), dental decay and rarely by orofacial cracks and pain, oral cancer and HIV lesions (Jin et al. 2016). Cavities and tooth decay are complex multifactorial infections with tooth plaque as their key reason (Shafei et al. 2012). Dental caries is ubiquitous process caused by a confined chemical

dissipation of the tooth surface prompted by acid produced due to the dental plaque exposed frequently to sugars (Fejerskov et al. 2015). Ulatrophia includes gingivitis and periodontitis are inflammatory infections of dental biofilm. Vital risk factor related to the inappropriate and damaging inflammatory response is the accretion of dental biofilm under the gingival margin (Chapple et al. 2015; Jepsen et al. 2017). Dental caries and periodontitis are caused by the imbalance of the dental biofilm (Teanpaisan et al. 2017). Dental plaque or biofilms are formed by the adherence and pellicle formation (thick layer of glycoproteins) on to the tooth surface that is instantly formed on clean surface. Ultimately, the sticky and soft deposition that is apparently observed on tooth surfaces clinically termed as dental

plaque (Manjia et al. 2018). Removal of bacterial plaque biofilms remains a challenge for health care centers. Oral microflora approximately comprises of 700 microbial species and 19,000 genotypes (phylotypes) (Keijser et al. 2008). In natural microflora of human body, the most diverse range of microbes present in oral cavity that deal with more than 1000 species (Manji et al. 2018). Endogenous bacteria produce organic acid that decreases plaque pH, which leads to demineralization of the tooth surface (Kim et al. 2017). Various factors including suitable pH, temperature and moist environment enhances the bacterial growth (Marsh et al. 2011).

Periodontal diseases and tooth decay (caries) are not caused by specific pathogens. The mechanism of interaction between microbiota-environment and host is much more complex (Colombo and Tanner 2019). Worldwide prevalence of dental caries is high and untreated dental caries that effecting human beings frequently (GBD 2016). As per recent global estimations, untreated caries in dentine in primary teeth and in permanent teeth are 621 million in children and 2.4 billion in all age group population respectively (Kassebaum et al. 2015). Globally almost 743 million people are affected by severe periodontitis (Jepsen et al. 2017). Various types of bacteria can grow and persist in the periodontal diseases and gingival crevice (Newman et al. 2006). Supragingival and subgingival are the two types of plaque. Supragingival portion possess mostly gram-positive bacteria including *Streptococcus sanguinus*, *Streptococcus mutans*, *Streptococcus salivarius*, *Streptococcus mitis* and *Lactobacilli*. Gram negative bacteria (*Actinomyces comitans*, *Actinobacillus*, *Fusobacterium nucleatum*, *Porphyromon asgingivalis*, spirochetes such as *Treponema denticola* and *Prevotella intermedia* primarily resides in subgingival plaque (Aas et al. 2005; Paster et al. 2001).

Medicinal plants produce different kind of metabolites and antimicrobial components that can be effective for the treatment of various infections and also have long term beneficial health effects. Secondary metabolites effectively have therapeutic actions in humans. From beginning to till current era almost one third of the total plant species are used for the treatment of different diseases due to the presence of secondary metabolites (Agbor and Naidoo 2015). For the treatment of different oral diseases and bacterial resistance against antibiotics, an effective therapeutic agent (secondary metabolites) of plants is being used in developing countries all over the world. Plants based treatment is extremely effective, economical, easily available and safe for the immune-compromised patients as an alternative to other preventive measures (Palombo 2011). Herbal products have antiviral, antifungal, antibacterial and anti-inflammatory activities and have found its way into dental medicine (Karadaghloglu et al. 2019). Herbal products can inhibit the biofilm formation and reduce the microbial adhesion which is the main cause of dental plaque and caries (Barnes and Arnold 2018).

Walnut tree (*Juglan regia L.*) bark is a main source of Derum is one of the natural "chewing sticks" and also mainly used in cosmetics products as a dye and as a tooth brush (Kadum and Al-Waheb 2010). The family Juglandaceae, genus *Juglans* (family) includes numerous species and it is present worldwide. Walnut is its prominent member, constituting a substantial species of plants which

are commonly present in the moderate climate areas and cultivated at commercial scale in the Asia, western South America, United States and southern and central Europe (Pereira et al. 2007).

The particular characteristics of walnut plant made it as effective substitute to antibiotics, chemical preservatives and antioxidant activity. Recent research evaluated the antimicrobial activity of phenolic extracts of walnut as well (Sousa et al. 2006; Muzaffer and Paul 2018). Various extract of medicinal plants has been reported as inhibitor of dental biofilm by reducing the adherence of pathogens to the tooth surface, that's leads to the progression of dental decay (Jain et al. 2015).

Walnuts comprises of different polyunsaturated fatty acids, amino acids, phenolic compounds, dietary fatty acids and minerals (Kavuncuoglu et al. 2018). Folic acid, β -sitosterol, Gallic acid, ascorbic acid, regiolone and quercetin-3- α -L-arabinoside are present in stem bark. Several studies show the antioxidant, antifungal and antibacterial activity of this plant (Noumi et al. 2010, Zakavi et al. 2013).

The current experiment estimates the determination of phenolic acids of plant extract through High performance liquid chromatography (HPLC) and antibacterial effect against oral microorganisms associated with periodontitis and dental caries. Cytotoxicity of different solvent extracts was also examined to find the CC_{50} .

MATERIAL AND METHODS

Acquisition of Medicinal plants

Was attained from local source (Faisalabad city). Plants were verified from Botany Department University of Agriculture, Faisalabad, Pakistan.

Preparation of plant extract

Bark was dried under shade and grinded by a grinding mill. Total 20 gram of sample was extracted in deionized water 200ml, acetone 70% and methanol 70%, respectively (Sultana et al. 2007). Extract was filtered by using vacuum filtration assembly using Whatman No. 1 filter paper. Through rotary evaporator filtrate was dried and stored at 4°C for further use.

Bacterial Strains

Three bacterial strains including *S. mitis* (ATCC 6249), *S. mutans* (ATCC 25175) and *P. intermedia* (ATCC 25611) were procured from Institute of Microbiology, University of Agriculture, Faisalabad. Salivarius agar (Himedia, India) was used as selective medium for *S. mutans* and *S. mitis* at 37°C for 24 hrs under aerobic conditions. Basal agar (Himedia, India) was used for the anaerobic growth of *P. intermedia*, and incubated under anaerobic conditions for 72 hrs.

Bacterial inoculum Preparation

McFarland (0.5) standard was used for the adjustment of bacterial suspension through spectrophotometer.

Evaluation of extract as an antimicrobial agent

For the assessment of antimicrobial activity of different extracts of agar well diffusion method was used with minor modifications (Andualet et al. 2014). For the aerobic and

facultative anaerobic bacterial strains, Mueller Hinton agar and basal agar were used, respectively. Broth culture of bacterial strains standardized with McFarland standard and streaked with sterilized cotton swab on agar plates, following this after 30 min by using cork borer wells of 6mm diameter were made on each plate equidistantly. Each well poured with 50µL of the extract solution. Aerobic and facultative anaerobic bacterial plates were incubated aerobically and anaerobically at 37 °C for 24 hrs and 48 hrs, respectively. After the incubation, zone of inhibition was observed of each type of extract against all the bacterial strains that shows the antimicrobial potential of respective extract. Diameters of the zones were measured in millimeters and all the extracts were assessed in triplicate for the mean value.

Determination of minimum inhibitory concentration (MIC)

For the determination of the minimum inhibitory concentration of plant extract, microdilution method was adopted with minor modifications (Amanullah et al. 2018). Dimethyl sulfoxide (DMSO) 100ul was used as a solvent for dry plant extract. Mueller Hinton broth and anaerobic basal broth were serially diluted (twofold dilution) with plant extract in microtitration plate. Then in 96 well microtitration plate 100 ul standardized inoculum of bacterial strains were added and incubated at 37°C for 24 hrs. 40 µl of 0.2 mg/ml of P-iodonitrotetrazolium chloride (INT) was added as a bacterial growth indicator in all wells and incubated for 30 min at 37 °C. Hence the change in the color due to the INT was used to evaluate the MIC (Eloff 1998). The well with no color change in medium was confirmed as MIC of sample concentration.

Minimum bactericidal concentration (MBC)

It is the minimum concentration of extract at which the tested bacteria were killed. MBC of various extract determined by taking 10 ul from each well that have no visible growth and poured on the respective agar plates and results were evaluated after 24 hrs incubation at 37°C.

Cytotoxicity assay (MTT)

Microculture MTT method was followed for determination of cytotoxicity of plant extracts. Each well is harvested with $2.0\text{--}2.8 \times 10^5$ cells and inoculated in 24 well plates. Phosphate buffer saline was used for the washing of cells and after that inoculated with and without the extract. Each sample was inoculated two times. Then, the cells were treated within MTT (Artun et al, 2016) and incubated at 37°C for 48 hours, 20µl of 5 g/l MTT was added to all the well and incubated for 4 hrs. Formazan crystals were solubilized by addition of warm 200 ul DMSO after removal of MTT from each well. The results were examined at 490 nm wavelength. Control wells cultivated without drugs in same conditions. Data were calculated as percentage of inhibition by the following formula.

$$\text{Inhibition \%} = \left[100 - \left(\frac{\text{ODt}}{\text{ODs}} \right) \times 100 \right] \%$$

HPLC analysis of plants extracts

For the phytochemical analysis of plant extract HPLC method with minor changes was used (San and Yildirim 2010). Initially, mobile phase was goes through degasser DGU-12A, Shimadzu, Japan, for the removal of any kind of bubble. Qualitative and quantitative analysis of sample were done by comparison with standard curves. Different phenolic acids present in plant extract were estimated through suitable mathematical calculations.

Statistical analysis

Statistical Package for Social Sciences Version 17 (SPSS Inc. Chicago, U.S.A) was used for the data analysis.

RESULTS

Antibacterial Effects of Extracts

The various extracts of *J. regia* exhibited variable degree of antibacterial activity against oral pathogenic bacteria. All the plant extract tested with different concentrations against all bacterial pathogens. Methanolic extract showed highest significant results against all bacterial strains. Acetone and aqueous extracts observed as less effective as compared to methanolic extract. Tables (1, 2, 3) and Figures (1, 2, 3) explained the results respectively of *S. mitis*, *S. mutans* and *P. intermedia*.

Table 1: Antimicrobial activity *J. regia* aqueous, acetone and methanol extract against *S. mitis*

Concentration mg/ml	Aqueous (Zone of inhibition in mm)	Acetone (Zone of inhibition in mm)	Methanol (Zone of inhibition in mm)
10	10.00±0.58D	8.33±0.88C	8.33±0.33D
20	11.33±0.94D	8.67±0.67C	9.00±0.58D
40	13.00±1.41CD	9.67±0.88BC	10±1.15CD
80	16.00±1.63BC	10.33±0.67BC	12.67±0.33BC
160	20.00±0.58A	13.00±1.15B	15.00±1.41AB
Vancomycin	16.67±0.33AB	16.67±0.33A	16.67±0.33A
PBS	0±0E	0±0D	0±0E
Overall Mean	12.43±1.36	9.52±1.08	10.24±1.15

Values are mean diameter of inhibition zone in millimeter (mm) ± S.E of three replicate.

The Means sharing similar letter in a row or in a column are statistically non-significant ($P>0.05$). PBS; phosphate buffer saline, vancomycin (30 µg) was used as reference antibacterial compound.

Table: 2: Antimicrobial activity *J. regia* aqueous, acetone and methanol extract against *S. mutans*




Concentration mg/ml	Aqueous (Zone of inhibition in mm)	Acetone (Zone of inhibition in mm)	Methanol (Zone of inhibition in mm)
10	0±0D	0±0D	0±0D
20	8±0.58C	0±0D	8.33±0.88BC
40	9.67±0.88C	8.5±1.41C	10.67±0.88BC
80	12.67±0.67B	10.00±0.58B	11.67±1.2
160	17±0.58A	12±0.58B	13±1.41AB
Vancomycin	16.67±0.33A	16.67±0.33A	16.67±0.33A
PBS	0±0D	0±0D	0±0D
Overall Mean	9.14±1.47	6.67±1.41	8.62±1.35

Values are mean diameter of inhibition zone in millimeter (mm) ± S.E of three replicate.

Table 3: Antimicrobial activity *J. regia* aqueous, acetone and methanol extract against *P. intermedia*

Concentration mg/ml	Aqueous (Zone of inhibition in mm)	Methanol (Zone of inhibition in mm)	Acetone (Zone of inhibition in mm)
10	0±0C	8.33±0.33C	0±0D
20	0±0C	8.33±0.33C	0±0D
40	0±0C	10.33±1.2C	0±0D
80	0±0C	13.67±0.33AB	8.33±0.33B
160	8±0.58B	15.67±0.33A	10.33±0.88AB
Clindamycin	11±0.58A	11±0.58BC	11±0.58A
PBS	0±0C	0±0D	0±0D
Overall Mean	2.71±0.98	9.62±1.05	4.24±1.12

Values are mean diameter of inhibition zone in millimeter (mm) ± S.E of three replicate.

**Fig. 1:** Graphic presentation of antimicrobial activity of *J. regia* aqueous, acetone and methanol extract against *S. mitis*.**Fig. 2:** Graphic presentation of antimicrobial activity of *J. regia* aqueous, acetone and methanol extract against *S. mutans*.**Fig. 3:** Graphic presentation of antimicrobial activity of *J. regia* aqueous, acetone and methanol extract against *P. intermedia*.

The Means sharing similar letter in a row or in a column are statistically non-significant ($P>0.05$). PBS; phosphate


buffer saline, vancomycin (30 µg) was used as reference antibacterial compound.

The means sharing similar letter in a row or in a column are statistically non-significant ($P>0.05$) PBS; phosphate buffer saline, Clindamycin (30 µg) was used as reference antibacterial compound.

Minimum Bactericidal Concentration (MBC) and Minimum inhibitory concentration (MIC): MIC and MBC were examined by using different concentrations of all three extract of (Table 4 and Figure 4).

Table 4: Minimum inhibitory concentration (MIC mg/ml) of different solvent extracts of *Juglans regia* against *P. intermedia*, *S. mitis* and *S. mutans*


Microorganism	Aqueous	Acetone	Methanol
<i>Streptococcus mutans</i>	20.00±0.00b	40.00±0.00a	20.00±0.00b
<i>Streptococcus mitis</i>	5±0.00a	10.00±0.00b	10.00±0.00b
<i>Prevotella intermedia</i>	160.00±0.00c	80.00±0.00a	10.00±0.00b

**Fig. 4:** Graphic presentation of MIC of different extracts of *J. Regia* against bacteria.

The aqueous and methanol extracts showed significantly higher activity (20.00±0.00, 20.00±0.00mg/ml each) followed by acetone extract against *S. mutans*. The aqueous extract showed significantly higher activity against *S. mitis*. The methanol extract was significantly effective than aqueous and acetone extract against *P. intermedia*. Table 5 and Fig. 5 showed MBC values of tested bacteria. The MBC of all extracts was higher than MIC against *S. mitis*. The MBC of methanolic, aqueous and acetone were 20.00±0.00 mg/ml 320.00±0.00, 160.00±0.00 correspondingly than MIC against *P. intermedia*.

Table 5: Minimum bactericidal concentration (MBC mg/ml) of different solvent extracts of *Juglans regia* against *P. intermedia*, *S. mitis* and *S. Mutans*

Microorganism	Aqueous	Acetone	Methanol
<i>Streptococcus mutans</i>	20.00±0.00b	40.00±0.00a	40.00±0.00a
<i>Streptococcus mitis</i>	10.00±0.00c	20.00±0.00a	20.00±0.00a
<i>Prevotella intermedia</i>	320.00±0.00c	160.00±0.00a	20.00±0.00b

**Fig. 5:** Graphic presentation of MBC of different extracts of *J. Regia* against bacteria.

Cytotoxicity assay

Dose dependent cytotoxicity of all three solvent extracts of were observed to Vero cells. Cell growth was suppressed with the higher concentrations of extracts inhibit the more cells. Table (6, 7, 8) showed the inhibitory percentage of methanolic, acetone and aqueous extract respectively.

HPLC analysis of *A. nilotica*

For the determination of phenolic acids in all three extracts of, HPLC was performed and methanolic extract has higher concentration of phenolic acids than aqueous and acetone extracts. Concentration of phenolic acids of all three extracts given in Table 10, Fig. 7.

Cellular Cytotoxicity (CC₅₀)

Regression analysis was used to calculate the cellular cytotoxicity (CC₅₀) from the observed data of inhibition dose response. The methanolic, acetone and aqueous extract showed CC₅₀ results as 278.43±29.77 µg/ml, 632.78±24.38 µg/ml and 929.65±47.48 µg/ml, respectively. Aqueous extract shows significantly higher CC₅₀ than acetone and methanolic extract. Table 9 and Fig. 6 explain the CC₅₀ results.

Table 6: Percentage inhibition of Vero cells by *J. regia* (aqueous)

Concentration µg/ml	Absorbance	%age inhibition	t vs control
Control	0.97±0.06	-	-
156.25	0.66±0.03	32.45±3.03	10.491
312.5	0.49±0.04	49.45±4.53	10.807
625	0.4±0.04	58.45±4.17	13.928
1250	0.35±0.03	64.39±3.5	18.314
2500	0.24±0.03	74.83±2.99	24.927

Assay were performed in triplicate Mean±SE.

Table 7: Percentage inhibition of Vero cells by *J. regia* (methanol)

Concentration µg/ml	Absorbance	%age inhibition	t vs control
Control	0.97±0.06	-	-
156.25	0.63±0.02	35.47±1.67	20.813
312.5	0.51±0.01	47.6±0.85	55.652
625	0.38±0.02	61.2±2.35	25.876
1250	0.34±0.02	64.9±2.33	27.755
2500	0.28±0.02	71.53±2.44	29.223

Assay were performed in triplicate Mean±SE.

Table 8: Percentage inhibition of Vero cells by *J. regia* (acetone)

Concentration µg/ml	Absorbance	%age inhibition	t vs control
Control	0.97±0.06	-	-
156.25	0.63±0.01	34.99±1.19	28.776
312.5	0.49±0.01	49.66±0.62	78.693
625	0.37±0.01	61.4±1.49	41.027
1250	0.36±0.01	62.98±0.54	116.734
2500	0.28±0.02	71.5±2.15	33.102

Assay were performed in triplicate Mean±SE.

Table 10: Phenolic of different solvents of *J. regia* (ppm)

Solvent	Caffeic Acid (ppm)	Vanillic Acid (ppm)	4-Hydroxy 3 Methoxy benzoic Acid (ppm)	Syringic Acid (ppm)	M-Comeric Acid (ppm)	Cromatotropi c Acid (ppm)	Gallic Acid (ppm)	Chlorogenic Acid (ppm)	P-Comeric Acid (ppm)	Ferulic Acid (ppm)	Quercetin
Acetone	0.00±0.00C	20.80±0.15B	69.50±0.36A	0.00±0.00B	0.00±0.00	0.00±0.00C	0.00±0.00C	0.00±0.00B	0.00±0.00	0.00±0.0	16.51±0.17B
Aqueous	3.80±0.15B	10.70±0.36C	0.00±0.00B	19.90±0.06A	0.00±0.00	42.30±0.35A	5.90±0.10B	0.00±0.00B	0.00±0.00	0.00±0.0	7.03±0.08C
methanol	14.50±0.32A	30.02±0.65A	0.00±0.00B	0.00±0.00B	2.90±0.21A	36.70±0.53B	15.60±0.26A	13.26±0.18A	2.80±0.25A	0.00±0.0	24.84±0.23A

Table 9: CC₅₀ µg/ml of different extracts of *J. regia*

Aqueous	568.35 ± 25.33a
Acetone	481.05 ± 79.68c
Methanol	492.98 ± 87.38b




Fig. 6: Graphic presentation of CC₅₀ of different *J. regia* extracts.




Fig. 7: Graphic presentation of phenolic acid in different extracts of *J. regia*.

DISCUSSION

Globally among many of major health problems oral diseases are still a major concern (Dash et al. 2014). Other than dental caries and periodontal diseases some other including oral cavity lesions, cancer of oral cavity and throat tissues also have great importance (Petersen 2003). Oral hygiene has great impact in general wellbeing of humans. Quality of life and whole-body systems other than craniofacial complex are deeply concerned with oral health. Treatment for the oral diseases and caries is much expensive in many countries (Sheiham 2005). For the treatment of oral diseases many natural and herbal remedies are being practiced since long time. Herbal and medical plants are very effective against many diseases. Researchers and health care authorities still working on many natural products against oral pathogens (Palombo 2011). Secondary metabolites of plants including phenolic compounds, are globally present in different vegetables, fruits and dietary products. Mainly phenolics are divided into three main groups including phenolic acids, flavonoids and tannins (Borges et al. 2013). The current study was conducted to evaluate the antibacterial activity of different extracts (methanolic, aqueous acetone) extract of *against*

oral pathogenic bacteria linked with periodontal diseases and caries.

The leaves are an easy to get remedy for prevention of oral diseases (Sharafati-Chaleshtori et al. 2011). Phenolic acids and flavonoids are extensively used in various pharmaceutical products due to their antioxidant activity. (Zheng and Wang 2001). Moreover, antibacterial potential of phenolic compounds has been already confirmed by different researchers worldwide because of this property phenolic extract of medicinal plants are considered as alternative to the antibiotics (Oliveira et al. 2008). Walnut is considered as an excellent source of antimicrobial compounds. Against *S. mutans* aqueous extract of evaluated statistically significant antimicrobial activity (17.00 ± 0.58 mm) (MIC 20 mg/ml) followed by methanol and acetone extracts. While against *S. mitis*, aqueous extract (20.00 ± 0.58 mm) was significantly effective too that followed by methanol and acetone, but against *P. intermedia*, methanol extract showed significant results (15.67 ± 0.33 mm). that followed by acetone and aqueous extract. The findings of current study were in accordance with the results of (Darmani et al. 2006; Pereira et al. 2007) which reported that aqueous extract of showed significant growth inhibition of oral pathogenic bacteria *Lactobacillus casei*, *S. mutans*, *S. salivarius* and *Actinomyces Viscosus*. The antibacterial including gram positive and gram negative, and antifungal potential of were evaluated by Alkhawajah (1997). Gram-negative bacteria have outer membrane that is made up of lipopolysaccharides covering, it can prevent hydrophobic compounds to diffuse into the cell but phytochemicals can pass easily through gram positive bacterial cell wall and also disturb active transport that results in coagulation of cells (Burt 2004). The MIC range from 20–40 mg/ml against *S. mutans*. MIC of Aqueous and methanol extract of was low (20 mg/ml) as compared to acetone extract that 40 mg/ml. The lower MIC was an indication of effectiveness of the extract. The MIC against *S. mitis* was 5.00 mg/ml of aqueous followed by acetone and methanol which was 10 mg/ml of each extract. The MIC against *P. intermedia* was 10 mg/ml of methanol followed by acetone 80 mg/ml than aqueous 160 mg/ml. the MBC value of three solvents were similar to MIC against *S. mutans*. MBC value of extracts were higher than the MIC value against *S. mitis* and *P. intermedia*. The aqueous and acetone extract of were tested against salivary flora as an antibacterial agent by (Deshpande et al. 2011; Jagtap and Karkera 2000). Various extract were also examined against Vero cells to estimate CC_{50} . Aqueous extract showed significantly higher (568.35 ± 25.33 μ g/ml) CC_{50} than the methanolic (490.98 ± 25.33 μ g/ml) and acetone (481.05 ± 79.68 μ g/ml) extract. The percentage of inhibition of cells were determined by dose response curve. Various extracts of showed no statistically significant difference between percentage inhibitions of cells. Ethanol extract of responded differently with different cell lines. The CC_{50} of ethanol extract was 100 μ g mL⁻¹. The different solvents showed divergent results that was in accordance with the finding of Chaieb et al (2013). The HPLC analysis of various extracts of were quite variable. Methanolic extract of has highest phenolic acids followed by aqueous and acetone extract. The findings of our HPLC results conform with the previous finding which reported

that extraction efficiency of phenolic compounds depends on solvent used. Phenolic compounds were found to be high in number in methanolic extract as compared to other solvents (Cosmulescu et al. 2010) and Pereira et al. (2007) reported that isolation and identification of ten phenolic compounds from walnut leaves. The phenols and flavonoids of walnut leaves are likely responsible for the antibacterial activities of the extract derived from (Stampar et al. 2006). Cueva et al. (2010) reported that phenolic acids has good inhibitory effect on intestinal bacteria and microbes in comparison with (-) epicatechin and (+) - catechin.

Conclusion

Methanol proved to be the best solvent for extraction of phenolic acids and its extract is most effective against pathogens under study. On the basis of CC_{50} , aqueous extract is the safest among the three solvent extracts to be used for therapeutic purposes. The results of present study suggest that *J. regia* may be used alone or in combination with other plant extracts in mouth wash or tooth paste to prevent oral diseases.

DECLARATIONS

Funding: This research work was Not funded by a national or international funding agency.

Acknowledgement: The authors appreciate the laboratory staff for their help.

Conflicts of Interest: Authors have no conflict of interest.

Availability of Data and Material: All the data is included in this study and can be obtained from corresponding author.

Ethics Statement: All the research procedures were conducted according to guidelines regarding use of animal for research purpose.

Authors Contribution: All the authors contributed equally.

Generative AI Statement: The authors declare that no Gen AI/Deep Seek was used in the writing/creation of this manuscript. **Publisher's Note:** All claims stated in this article are exclusively those of the authors and do not necessarily represent those of their affiliated organizations or those of the publisher, the editors, and the reviewers. Any product that may be evaluated/assessed in this article or claimed by its manufacturer is not guaranteed or endorsed by the publisher/editors.

Publisher's Note: All claims stated in this article are exclusively those of the authors and do not necessarily represent those of their affiliated organizations or those of the publisher, the editors, and the reviewers. Any product that may be evaluated/assessed in this article or claimed by its manufacturer is not guaranteed or endorsed by the publisher/editors.

REFERENCES

- Aas JA, Paster BJ, Stokes LN, Olsen I and Dewhirst FE 2005. Defining the normal bacterial flora of the oral cavity. *Journal of Clinical Microbiology* 43(11): 5721-5732.
- Agbor MA and Naidoo S 2015. Ethnomedicinal plants used by traditional healers to treat oral health problems in Cameroon. *Evidence-Based Complementary and Alternative Medicine* 649832: 1-10.
- Alkhawajah AM 1997. Studies on the antimicrobial activity of *Juglans regia*. *American Journal of Chinese Medicine* 25(2): 175-180.
- Amanullah M, Sadatullah S, Yaseen SM and Dawasaz AA 2018. Use of herbal extract from *Artemisia herba-alba* (Shih) in pharmaceutical preparations for dental hygiene. *Saudi Pharmaceutical Journal* 26(6): 822-828.
- Andualem G, Umar S and Getnet F 2014. Antimicrobial and phytochemical screening of methanol extracts of three medicinal plants in Ethiopia. *Advances in Biological Research* 8(3): 101-106.
- Artun FT, Karagoz A, Ozcan G, Melikoglu G, Anil S, Kultur S and Sutlupinar N 2016. In vitro anticancer and cytotoxic activities of some plant extracts on HeLa and Vero cell lines. *Journal of B.U.ON.* 21(3): 720-725.
- Barnes NA and Arnold PT 2018. Investigation of antibacterial effects of Southern Appalachian plants: *Phytolacca americana* and *Juglans nigra* on selected bacteria. *Georgia Journal of Science* 76(2): 45.
- Borges A, Ferreira C, Saavedra MJ and Simoes M 2013. Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. *Microbial Drug Resistance* 19(4): 256-265.
- Burt S 2004. Essential oils: their antibacterial properties and potential applications in foods—a review. *International Journal of Food Microbiology* 94(3): 223-253.
- Chaieb K, Kouidhi B, Slama RB, Fdhila K, Zmantar T and Bakhrouf A 2013. Cytotoxicity, antibacterial, antioxidant, and antibiofilm properties of Tunisian *Juglans regia* bark extract. *Journal of Herbs, Spices & Medicinal Plants* 19(2): 168-179.
- Chapple IL, Van der Weijden F, Doerfer C, Herrera D, Shapira L, Polak D, Madianos P, Louropoulou A, Machtei E, Donos N, Greenwell H, Van Winkelhoff AJ, Eren Kuru B, Arweiler N, Teughels W, Aïmetti M, Molina A, Montero E and Graziani F 2015. Primary prevention of periodontitis: managing gingivitis. *Journal of Clinical Periodontology* 42(S16): S71-S76.
- Colombo APV and Tanner ACR 2019. The role of bacterial biofilms in dental caries and periodontal and peri-implant diseases: a historical perspective. *Journal of Dental Research* 98(4): 373-385.
- Conway DI, McMahon AD, Robertson AD and Macpherson LMD 2013. Dental epidemiology. In: *Handbook of Epidemiology*, Ahrens W and Pigeot I (eds). Springer Scientific, New York.
- Cosmulescu S, Botu M, Trandafir I, Baciua A, Achim G and Gruia M 2010. Phenolics of green husk in mature walnut fruits. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 38: 53-56.
- Cueva C, Moreno-Arribas MV, Martín-Alvarez PJ, Bills G, Vicente MF, Basilio A, Rivas CL, Requena T, Rodríguez JM and Bartolomé B 2010. Antimicrobial activity of phenolic acids against commensal, probiotic and pathogenic bacteria. *Research in Microbiology* 161(5): 372-382.
- Darmani H, Nusayr T and Al-Hiyasat AS 2006. Effects of extracts of miswak and derum on proliferation of Balb/C 3T3 fibroblasts and viability of cariogenic bacteria. *International Journal of Dental Hygiene* 4(2): 62-66.
- Dash TR, Singh N, Gupta D, Panwar E and Ramisetty S 2014. Role of medicinal herbs in oral health management. *International Journal of Dentistry and Medical Research* 1(4): 113-119.
- Deshpande RR, Kaleb AA, Ruikar AD, Panvalkar PS, Kulkarnia AA, Deshpande NR and Salvekar JP 2011. Antimicrobial activity of different extracts of *Juglans regia* L. against oral microflora. *International Journal of Pharmaceutical Sciences and Research* 3(1): 200-201.
- Eloff JN 1998. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plant extracts for bacteria. *Planta Medica* 64(8): 711-713.
- Fejerskov O, Nyvad B and Kidd E 2015. Dental caries, what is it? In: *Dental Caries: The Disease and its Clinical Management*, Fejerskov O, Nyvad B and Kidd E (eds), 3rd edition, Wiley Blackwell, Oxford.
- GBD 2015 Disease and Injury Incidence and Prevalence Collaborators 2016. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet* 388(10053): 1545-1602.
- Jagtap AG and Karkera SG 2000. Extracts of *Juglans regia* inhibits growth in vitro adherence, acid production and aggregation of *Streptococcus mutans*. *Journal of Pharmacy and Pharmacology* 52(2): 235-242.
- Jain I, Jain P, Bisht D, Sharma A, Srivastava B and Gupta N 2015. Use of traditional Indian plants in the inhibition of caries-causing bacteria - *Streptococcus mutans*. *Brazilian Dental Journal* 26(2): 110-115.
- Jepsen S, Blanco J, Buchalla W, Carvalho JC, Dietrich T, Dorfer C, Eaton KA, Figuero E, Frencken JE, Graziani F, Higham SM, Kocher T, Maltz M, OrtizVigon A, Schmoekel J, Sculean A, Tenuta LMA, van der Veen MH and Machiulskiene V 2017. Prevention and control of dental caries and periodontal diseases at individual and population level: consensus report of group 3 of joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *Journal of Clinical Periodontology* 44(S18): S85-S93.
- Jin LJ, Lamster IB, Greenspan JS, Pitts NB, Scully C and Warnakulasuriya S 2016. Global burden of oral diseases: emerging concepts, management and interplay with systemic health. *Oral Diseases* 22(7): 609-619.
- Kadum MS and Al-Waheb AM 2010. Effect of derum (*Juglans regia* L. bark) extracts on *Mutans Streptococci* in comparison to chlorhexidine gluconate (in vitro study). *Journal of Baghdad College of Dentistry* 22(4): 83-87.
- Karadaglioglu OI, Ulusoy N, Baser KHC, Hanoglu A and Sık I 2019. Antibacterial activities of herbal toothpastes combined with essential oils against *Streptococcus mutans*. *Pathogens* 8(4): 1-17.
- Kassebaum NJ, Bernabe E, Dahiya M, Bhandari B, Murray CJL and Marcenes W 2015. Global burden of untreated caries: a systematic review and meta-regression. *Journal of Dental Research* 94(5): 650-658.
- Kavuncuoglu H, Kavuncuoglu E, Karatas SM, Benli B, Sagdic O and Yalcin H 2018. Prediction of the antimicrobial activity of walnut (*Juglans regia* L.) kernel aqueous extracts using artificial neural network and multiple linear regression. *Journal of Microbiological Methods* 148: 78-86.
- Keijser BJ, Zaura E, Huse SM, van der Vossen JM, Schuren FH, Montijn RC, ten Cate JM and Crielaard W 2008. Pyrosequencing analysis of the oral microflora of healthy adults. *Journal of Dental Research* 87(11): 1016-1020.
- Kim Y, Ha M, Kwon H, Kim H and Choi Y 2017. Association between low blood lead levels and increased risk of dental caries in children: a cross-sectional study. *BMC Oral Health* 17(1): 1-7.
- Manji F, Dahlen G and Fejerskov O 2018. Caries and periodontitis: contesting the conventional wisdom on their aetiology. *Caries Research* 52(6): 548-564.

- Marsh PD, Moter A and Devine DA 2011. Dental plaque biofilms: communities, conflict and control. *Periodontology* 2000 55(1): 16-35.
- Muzaffer U and Paul VI 2018. Phytochemical analysis, in vitro antioxidant and antimicrobial activities of male flower of *Juglans regia* L. *International Journal of Food Properties* 21(1): 360-371.
- Newman JRS, Ghaemmaghami S, Ihmels J, Breslow DK, Noble M, DeRisi JL and Weissman JS 2006. Single-cell proteomic analysis of *S. cerevisiae* reveals the architecture of biological noise. *Nature* 441(7095): 840-846.
- Noumi E, Snoussi M, Hajlaoui H, Valentin E and Bakhrouf A 2010. Antifungal properties of *Salvadora persica* and *Juglans regia* L. extracts against oral *Candida* strains. *European Journal of Clinical Microbiology & Infectious Diseases* 29(1): 81-88.
- Oliveira I, Sousa A, Ferreira ICFR, Bento A, Estevinho L and Pereira JA 2008. Total phenols, antioxidant potential and antimicrobial activity of walnut (*Juglans regia* L.) green husks. *Food and Chemical Toxicology* 46(7): 2326-2331.
- Palombo EA 2011. Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evidence-Based Complementary and Alternative Medicine* 680354: 1-15.
- Paster BJ, Beches SK, Galvirin JI, Ericson RE, Lai CN, Levanos VA, Sahasrabudie A and Dewhirst FE 2001. Bacterial diversity in subgingival plaque. *Journal of Bacteriology* 183(12): 3770-3783.
- Pereira JA, Oliveira I, Sousa A, Valentao P, Andrade P, Ferreira I, Ferreres F, Bento A, Seabra R and Estevinho L 2007. Walnut (*Juglans regia* L.) leaves: phenolic compounds, antibacterial activity and antioxidant potential of different cultivars. *Food and Chemical Toxicology* 45: 2287-2295.
- Petersen PE 2003. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century- the approach of the WHO Global Oral Health Programme. *Community Dentistry and Oral Epidemiology* 31(1): 3-24.
- Shafiei Z, Shuhairi NN, Yap NMFS, Sibungkil CH and Latip J 2012. Antibacterial activity of *Myristica fragrans* against oral pathogens. *Evidence-Based Complementary and Alternative Medicine* 825362: 1-7.
- Sharafati-Chaleshtori R, Sharafati-chaleshtori F and Rafieian M 2011. Biological characterization of Iranian walnut (*Juglans regia*) leaves. *Turkish Journal of Biology* 35(5): 635-639.
- Sheiham A 2005. Oral health, general health and quality of life. *Bulletin of the World Health Organization* 83(9): 644-645.
- Sousa A, Ferreira ICFR, Calhelha R, Andrade PB, Valentao P, Seabra R, Estevinho L, Bento A and Pereira JA 2006. Phenolics and antimicrobial activity of traditional stoned table olives "alcaparra". *Bioorganic & Medicinal Chemistry* 14(24): 8533-8538.
- Stampar F, Solar A, Hudina M, Veberic R and Colaric M 2006. Traditional walnut liqueur - cocktail of phenolics. *Food Chemistry* 95(4): 627-631.
- Teanpaisan R, Kawsud P, Pahumunto N and Puripattanavong J 2017. Screening for antibacterial and antibiofilm activity in Thai medicinal plant extracts against oral microorganisms. *Journal of Traditional and Complementary Medicine* 7(2): 172-177.
- Zakavi F, Golpasand Hagh L, Daraeighadikolaei A, Farajzadeh Sheikh A, Daraeighadikolaei A and Leilavi Shooshtari Z 2013. Antibacterial effect of *Juglans regia* bark against oral pathologic bacteria. *International Journal of Dentistry* 854765: 1-5.
- Zheng W and Wang SY 2001. Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agricultural and Food Chemistry* 49(11): 5165-5170.