



Antibacterial Potential of Honeybee (*Apis mellifera*) Venom against *Pseudomonas aeruginosa*

Mukarma Mahmood^{1*}, Muhammad Moazam Jalees¹, Muhammad Waseem Abbas³, Waqas Ashraf¹, Abdullah Saghir Ahmad², Zeeshan Nawaz⁴, Qudratullah⁵, Shah Nawaz⁶ and Roshan Riaz⁷

¹Department of Microbiology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur.

²Department of Parasitology, Cholistan University of Veterinary and Animal Sciences, Bahawalpur.

³Department of Pathology, Quaide Azam Medical College Bahawalpur.

⁴Institute of Microbiology, Government College University Faisalabad.

⁵Department of Surgery, Cholistan University of Veterinary and Animal Sciences Bahawalpur.

⁶Department of Comparative Biomedical Sciences, School of Veterinary Medicine, Faculty of Health and Medical Sciences, University of Surrey, United Kingdom,

⁷Department of Animal Nutrition and Nutritional Disease, Faculty of Veterinary Medicine, Kafkas University, Kars, Türkiye

*Correspondence: mukarmamahmood21@gmail.com

ARTICLE INFO

ARTICLE HISTORY: CVJ-25-127

Received: 21 November 2025

Revised: 23 December 2025

Accepted: 25 January 2026

Published online: 05 February 2026

Key words:

Antimicrobial activity

Apis mellifera

Bee venom

MIC

Pseudomonas aeruginosa

Zone of inhibition

ABSTRACT

The study investigated the antibacterial properties of honeybee venom against *P. aeruginosa*. Honeybee venom contains antibacterial compounds that can help to treat lung and urinary tract infections caused by the Gram-negative bacterium *P. aeruginosa*. A study involving 264 clinical samples from urine, sputum, and pus found that venom had potent antibacterial properties against *P. aeruginosa*. The study found that venom preparation techniques could affect antibiotic efficacy and warranted further research. The most common source of samples was pus, and wound infections, abscesses, and post-operative infections were common diagnoses. When PCR was done, a particular primer was found. A distinct band on the gel at 396 and 1600 base pairs (bp) indicates the expected size of the amplicon for the Exo A and 16S rRNA genes. The study suggests that venom can be used in combination with other drugs to combat *P. aeruginosa* and could be beneficial for treating infections caused by bacterial activity.

To Cite This Article: Mahmood M, Jalees MM, Abbas MW, Ashraf W, Ahmad AS, Nawaz Z, Qudratullah, Nawaz S, and Riaz R, 2026. Antibacterial potential of honeybee (*Apis mellifera*) venom against *Pseudomonas aeruginosa*. Continental Vet J, 6(1): 126-130. <http://dx.doi.org/10.71081/cvj/2026.067>

INTRODUCTION

The demand for natural medicines as alternatives to synthetic antibiotics is increasing due to financial constraints and a desire for improved health outcomes (Veetilvalappil et al., 2022). However, overuse, misuse, and incomplete drug regimens have led to bacterial resistance (Rabea et al., 2025; Ijaz et al., 2019). *Pseudomonas aeruginosa*, a Gram-negative bacterium, is a significant public health concern due to its rapid development of multidrug resistance (Fernández et al., 2023).

Honeybee venom, a complex mixture of bioactive peptides and enzymes, has broad-spectrum antibacterial properties. Antibiotic resistance is a growing global health concern, causing increased morbidity and mortality rates (Awad et al., 2025). Natural products like bee venom are being investigated as alternative therapies to combat resistant

bacterial strains (Uddin et al., 2016). This study will use conventional microbiological methods and PCR for precise identification of *P. aeruginosa*, enhancing specificity and reliability in confirming isolate presence (Mesaros et al., 2007). Bee venom's antimicrobial, anti-inflammatory, and immunomodulatory properties may increase host immune responses and reduce infection severity (Aurongzeb et al., 2021). Venom-derived medications are recognized for their superior bioactivity, specificity, and stability compared to synthetic alternatives (Reynolds et al., 2021). Honeybee venom (BV) contains biologically active compounds like peptides and enzymes, which have antibacterial, anti-inflammatory, and cytotoxic activities (Iorizzo et al., 2022). Melittin, the major component of BV, is particularly potent against Gram-positive and some Gram-negative bacteria, disrupting cell membranes and enhancing antibiotic effects. BV has historically been used as a supplemental medication for various ailments (Zolfagharian et al., 2016). Toxins in

venoms can be beneficial as therapeutic agents, drug design templates, and pharmacological instruments in drug development (Leandro et al., 2025). This study examined the antibacterial properties of BV and gentamicin against specific bacteria strains (Muller et al., 2007; Alsayeqh, 2025). *P. aeruginosa*, a major cause of nosocomial infections, is often linked to medical devices and hospital environments, with resistance mechanisms like biofilm formation complicating treatment strategies (El-Seedi et al., 2020).

MATERIALS AND METHODS

Collection and Preparation of Sample

Samples were collected from Victoria Hospital Bahawalpur, South Punjab, using sterile syringes or swabs (Ahsan et al., 2024). Hand extraction was used to extract HB venom from healthy *A. mellifera* populations. The venom was dried, cut, and stored at -20°C. The venom was then weighed, documented, and packed in jars. Two types of bee venom were evaluated for their antimicrobial effects on *Pseudomonas aeruginosa*, including hand-extracted and lyophilized forms, using controlled stimulation techniques.

Isolation of *Pseudomonas aeruginosa*

The study involved preparing MacConkey agar plates and Cetrimide agar plates (selective media) and incubating specimens for one to two days to isolate *P. aeruginosa* strains. Morphological and physiological tests were used to identify them. 264 *Pseudomonas* isolates were collected from clinical samples (Abbas et al., 2022).

Bacterial Straining and Culturing on different Agars

MacConkey agar medium from Oxoid company was used for identifying and isolating *P. aeruginosa*, sterilized and autoclaved. Colonies were separated and incubated at 35°C for 24 to 48 hours. Samples were streaked on Cetrimide nalidixic acid agar using sterile cotton swabs for eight hours, then incubated at 37 degrees Celsius for 24 hours.

Transfer a small amount of suspended culture to a microscope slide, cover with a thin layer, and gently flame-dry or air-dry to avoid ring formations and overheating. A colony from MacConkey and Cetrimide Agar is fixed with crystal violet, then washed with tap water. Gram's iodine is used as a mordant, then cleaned with water. The slide is then cleaned with 95% alcohol, ethanol-acetone decolorizers, and safranin. The slide is then air-dried, and the counterstain is removed. The process involves rinsing, washing, and air-drying the slide to ensure the integrity of the culture (Rasool et al., 2016).




Fig 1: Colorless colonies on MacConkey agar.

Subculture on Cetrimide Agar

P. aeruginosa colonies were formed on Cetrimide agar, producing two colors: pyoverdine and pyocyanin, and all 28 positive cultures were found on Cetrimide agar (Nader et al., 2021).

Production of pyoverdine pigment




Fig 2: Results showing production of pyoverdine pigment on cetrimide agar.

Biochemical Testing

The study conducted biochemical assays on bacterial isolates, revealing that all tested positive for catalase, an enzyme that converts hydrogen peroxide into H₂O and O₂, and oxidase, a dark purple color indicating enzyme formation within five to ten seconds (Al-Bayati et al., 2021).

Production of pyocyanin pigment




Fig 3: Results showing production of pyocyanin pigment on cetrimide agar.

Purple color showing positive results




Fig 4: Colony smeared on filter paper by using Oxidase enzyme, purple hue emerged.

Oxidase test: This assay confirmed the presence of the cytochrome c oxidase enzyme, crucial in ETC, using a sterile wooden stick, filter paper, bacterial culture, petri plates, and 1% tetramethyl p-phenylenediamine dihydrochloride. Safety precautions were followed, including using fresh cultures and avoiding metal loops (Wellinghausen et al., 2025).

Catalase test: This test requires sterile swabs, glass slides, hydrogen peroxide, and bacterial culture. Fresh cultures are inoculated, and safety precautions are taken (Su et al., 2018).




Fig 5: Bubbles showing positivity in results of on glass slide.




Fig 6: Green color shows negative results and blue color shows positive results.

Simmons citrate test: It involved dissolved citrate agar in distilled water, poured into angled tubes, and inserted into a bacterial needle. The test showed a blue hue, indicating alkalinity, after a full day of tilting (Su et al., 2018).

PCR

The PCR technique uses specific primers for target genes to identify amplified DNA. It involves denaturation, annealing, extension, and amplification at 95°C and 72°C (Liu et al., 2023). Results are detected by electrophoresis in an agarose gel and examined with an ultraviolet transilluminator. The size of the amplicon is determined using a 5.5% agarose gel (Gariyban et al., 2013; Basit et al., 2021).




Fig 7: Agarose gel electrophoresis of a PCR amplification targeting the Exo A gene in *Pseudomonas aeruginosa*.




Fig 8: Agarose gel electrophoresis of a PCR amplification targeting the 16S rRNA gene in *Pseudomonas aeruginosa*.

Antibacterial Assay

Disk diffusion method used to study honey bee venom. Filter paper disks moistened in the test material are used in this manner. The venom of honey bee is put on a bacterially

contaminated agar plate. The zone of inhibition exhibits increased antibacterial activity after incubation (Abdulhaq et al., 2020; Shahid et al., 2023).

MIC

The MIC of bee venom was determined using the broth microdilution method, using 96-well microplates. The bacterial inoculum concentration in each well was 5x10⁶ CFU/ml. Ten dilutions were made, with a 1000 µg/mL stock solution made by mixing bee venom with nutrient broth. Three wells were used as background and growth controls. The MIC value was calculated using a spectrophotometer. Bacterial strains were cultivated in broth media, and the MIC of venom inhibited over 90% of the strains (El Sherry et al., 2025).

RESULTS AND DISCUSSION

Antibacterial activities of Bee venom revealed the significances opposed to *P. aeruginosa* (Gellatly et al., 2013 et al., 2013). There were different zones observed, like at 75 µg/mL 18 mm (the largest zone of inhibition), at 50 µg/mL 16 mm and at 25 µg/mL 14 mm (smaller zone) were observed in lyophilized venom. In Hand extracted venom, at 75 µg/mL 16mm (the largest zone), at 50 µg/mL 15mm and at 25 µg/mL 12mm (smaller zone) were observed (Moradali et al., 2007).

The study found *P. aeruginosa* in 10.6% of clinical samples, with the highest positivity rates in post-surgical infections, urinary tract infections, and pyelonephritis (Hanock et al., 2013). No *P. aeruginosa* isolates were found in pneumonia cases. PCR amplification confirmed culture-based identifications of *P. aeruginosa*, confirming its role in virulence and conserved sequence among clinical isolates, making it a reliable genetic marker (Lavenir et al., 2007).

The study found differential antimicrobial activity between lyophilized and hand-extracted venom, suggesting lyophilization may preserve bioactive compounds (Poirel et al., 2020). The higher prevalence of *P. aeruginosa* in hospital settings highlights the need for enhanced infection control measures (Poirel et al., 2010).

DECLARATIONS

Conflict of Interest: The authors declare no conflict of interest.

Funding: This research was not funded by any international or national funding agency.

Acknowledgement: The authors acknowledge the help provided by staff of department Microbiology of Cholistan University of Veterinary and Animal Sciences, Bahawalpur, Pakistan.

Authors' Contribution

M Jalees supervised and designed the study and M. Mahmood collected the bees for hand extraction method. Abdullah Saghir Ahmad, Muhammad Waseem Abbas, Waqas Ashraf, Zeeshan Nawaz, Qudratullah, Shah Nawaz and Roshan Riaz actively participated in preparation of manuscript, editing and finalization. Each author helped

write the manuscript, checked the finished version, and gave their approval for submission.

Data Availability: All the data is included in this paper and may be obtained from corresponding author.

Publisher's Note: All claims stated in this article are exclusively those of the authors and do not necessarily represent those of their affiliated organizations or those of the publisher, the editors, and the reviewers. Any product that may be evaluated/assessed in this article or claimed by its manufacturer is not guaranteed or endorsed by the publisher/editors.

REFERENCES

- Al-Bayati SS, Al-Ahmer SD, Shami AMM and Al-Azawi AH, 2021. Isolation and identification of *Pseudomonas aeruginosa* from clinical samples. *Biochemical Cell Archive*, 21: 3931-3935.
- Abbas R, Nawaz Z, Siddique AB, Aslam R, Rafique A, Zahoor MA, Qamar MU, Ahmad MZ, Jalees MM, Qasim M and Alsayeqh AF, 2022. Molecular detection of biofilm production among multidrug resistant isolates of *Pseudomonas aeruginosa* from meat samples. *Pakistan Veterinary Journal*, 42:505-510.
- Adulhaq N, Nawaz Z, Zahoor MA and Siddique A, 2020. Association of biofilm formation with multi drug resistance in clinical isolates of *Pseudomonas aeruginosa*. *EXCLI Journal*, 19:201-208.
- Ahsan H, Ayub M, Gul M, Qureshi A, Asfour HZ, Bilal HM, Azeem M, Wahid A, Ali N, Naveed R, Shabir M, Rajeh N, Mammadov A and Siddique AB, 2024. Efficacy of silver oxide nanoparticles against multi-drug resistant *Pseudomonas aeruginosa* and methicillin-resistant *Staphylococcus aureus* in burn wound infections. *Asian Journal of Agriculture and Biology*, 2024:2024096.
- Alsayeqh AF, 2025. Antibiotic resistance and emerging alternatives for controlling foodborne pathogens. *Pakistan Veterinary Journal*, 45:499-514.
- Aurongzeb M and Kamran M, 2011. Antimicrobial properties of natural honey: a review of literature. *J. Biochem. Mol. Biol*, 44:118-124.
- Awad EH, Arafa W, Ali HR, Barakat OS and Ahmed MN 2025. Unveiling the anti-biofilm potential of bee venom against multi-drug resistant human pathogenic bacteria and fungi: perspectives into the efficacy and Possible mechanisms. *Microbial Pathogenesis*, 200:107358.
- Basit M, Siddique AB, Aslam B, Zahoor MA, Hussain R, Ulhaq M, 2021. Distribution and antimicrobial susceptibility profile of bacterial and fungal pathogens from burn wounds in hospitalized patients. *Journal of Pakistan Medical Association*, 71:916-920
- El-Seedi H, Abd El-Wahed A, Yosri N, Musharraf SG, Chen L, Moustafa M, Zou X, Al-Mousawi S, Guo Z, Khatib A and Khalifa S, 2020. Antimicrobial Properties of *Apis mellifera*'s Bee Venom. *Toxins*, 12(7).
- EL-Sherry S, Nasr A and Zayed G, 2025. Triton X-100 -coated toltrazuril nanoparticles: a promising anti-coccidial formula for effective treatment of poultry coccidiosis. *Pakistan Veterinary Journal*, 45:1876-1884.
- Fernández L and Hancock RE, 2012. Adaptive and mutational resistance: Role of porins and efflux pumps in drug resistance. *Clinical Microbiology Reviews*, 25:661-681.
- Garibyan L, 2013. Research Techniques Made Simple: Polymerase Chain Reaction (PCR). *Journal of Investigative Dermatology*, 133:1-4.
- Gellatly SL and Hancock REW, 2013. *Pseudomonas aeruginosa*: New insights into pathogenesis and host defenses. *Pathogens and Disease*, 67:159-173.
- Hancock and Fernández LRE, 2012. Adaptive and mutational resistance: Role of porins and efflux pumps in drug resistance. *Clinical Microbiology Reviews*, 25:661-681.
- Han S, Zaniewski RP, Marr ES, Lacey BM, Tomaras AP, Evdokimov A, Miller JR and Shanmugasundaram V, 2010. Structural basis for effectiveness of siderophore-conjugated monocarbams against clinically relevant strains of *Pseudomonas aeruginosa*. *Proceedings of the National Academy of Sciences*, 107:22002-22007.
- Ijaz M, Siddique AB, Rasool MH and Shafique M, 2019. Frequency of multi drug resistant *Pseudomonas aeruginosa* in different wound types of hospitalized patients. *Pakistan Journal of Pharmaceutical Sciences*, 32:865-870.
- Iorizzo M, Letizia F, Ganassi S, Testa B, Petrarca S, Albanese G, Di Criscio D, and De Cristofaro A, 2022. Functional Properties and Antimicrobial Activity from Lactic Acid Bacteria as Resources to Improve the Health and Welfare of Honey Bees. *Insects*, 13:308.
- Lavenir R, Jocktane D, Laurent F, Nazaret S and Cournoyer B, 2007. Improved reliability of *Pseudomonas aeruginosa* PCR detection by the use of the *exoA* gene target. *Journal of Microbiological Methods*, 70: 20-29.
- Leandro LF, Mendes CA, Casemiro LA, Vinholis AHC, Cunha WR, Almeida RD and Martins CHG, 2015. Antimicrobial activity of apitoxin, melittin and phospholipase A2 of honey bee (*Apis mellifera*) venom against oral pathogens. *Anais Da Academia Brasileira de Ciências*, 87: 147-155.
- Liu D, Zhou Y, Li C, Li Y, Jiang Y, Huang Z, Liang J and Shu R, 2010. Denaturing gradient gel electrophoresis analysis with different primers of subgingival bacterial communities under mechanical debridement. *Microbiology and Immunology*, 54:702-706.
- Mesaros N, Nordmann P, Plésiat P, Roussel-Delvallez M, Van Eldere J, Glupczynski Y, Van Laethem Y, Jacobs F, Lebecque P, Malfroot A, Tulkens PM and Van Bambeke F, 2007. *Pseudomonas aeruginosa*: resistance and therapeutic options at the turn of the new millennium. *Clinical Microbiology and Infection: The Official Publication of the European Society of Clinical Microbiology and Infectious Diseases*, 13:560-578.
- Moradali MF, Ghods S and Rehm BHA, 2017. *Pseudomonas aeruginosa* lifestyle: A paradigm for adaptation, survival, and persistence. *Frontiers in Cellular and Infection Microbiology*, 7:39.
- Pang Z, Raudonis R, Glick BR, Lin TJ and Cheng Z, 2019. Antibiotic resistance in *Pseudomonas aeruginosa*: Mechanisms and alternative therapeutic strategies. *Biotechnology Advances*, 37:177-192.
- Muller JF, Stevens AM, Craig J and Love NG, 2007. Transcriptome Analysis Reveals that Multidrug Efflux Genes Are Upregulated to Protect *Pseudomonas aeruginosa* from Pentachlorophenol Stress. *Applied and Environmental Microbiology*, 73:4550-4558.
- Nader RA, Mackieh R, Wehbe R, El Obeid D, Sabatier JM, and Fajloun Z, 2021. Beehive Products as Antibacterial Agents: A Review. *Antibiotics*, 10:717.
- Poirer L, Jayol A and Nordmann P, 2020. Polymyxins: Antibacterial activity, susceptibility testing, and resistance mechanisms. *Clinical Microbiology Reviews*, 30:557-596.
- Rabea EY, Mahmoud ED, Mohamed NK, Ansary ER, Alrouby MR, Shehata RR, Mokhtar YY, Arullampalam P, Hegazy AM, Al-Sabi A, and Abd El- Aziz TM, 2025. Potential of Venom-Derived Compounds for the Development of New Antimicrobial Agents. *Toxins*, 17:238.
- Rasool MH, Siddique AB, Saqalein M, Asghar MJ, Zahoor MA, Aslam B, Shafiq HB, Nisar MA, 2016. Occurrence and

- antibacterial susceptibility pattern of bacterial pathogens isolated from diarrheal patients in Pakistan. Saudi Medical Journal, 37:274-279.
- Reynolds CR, Altmann RA and Allen DN, 2021. The Problem of Bias in Psychological Assessment. Mastering Modern Psychological Testing, 573–613.
- Shahid M, Hussain R, Nawaz Z, Aslam B, Ahmad MZ, Siddique AB, Ahsan H, Fatima A, Khan I, Mustafa B, Iqbal, Al Syaadn KM and Shami A, 2023. Occurrence of Virulence genes among Methicillin Resistant Staphylococcus aureus isolated from subclinical bovine mastitis. ACS Omega, 8:38111-38117.
- Su SS, Lae KZW and Ngwe H, 2018. Isolation and Identification of Pseudomonas aeruginosa from the Clinical Soil. University of Yangon Research Journal, 8:271- 275.
- Uddin MB, Lee BH, Nikapitiya C, Kim JH, Kim TH, Lee HC, Kim CG, Lee JS and Kim CJ, 2016. Inhibitory effects of bee venom and its components against viruses in vitro and in vivo. Journal of Microbiology, 54:853–866.
- Veetilvalappil VV, Manuel A, Aranjani JM, Tawale R and Koteswara A, 2022. Pathogenic arsenal of Pseudomonas aeruginosa: an update on virulence factors. Future Microbiology, 17:465–481.
- Wellingshausen N and Lander F, 2025. Rationale Infektionsdiagnostik in der Pädiatrie. Infektiologie Up2date, 1: 161-182.
- Zolfagharian H, Mohajeri M and Babaie M, 2016. Bee Venom (Apis Mellifera) an Effective Potential Alternative to Gentamicin for Specific Bacteria Strains: Bee Venom an Effective Potential for Bacteria. Journal of Pharmacopuncture, 19:225–230.