



Review Article

Phytomedicine efficacy and prospects in poultry; a new insight to old anthelmintic resistance

Saba Rashid^{1*}, Faisal Hafeez², Rehan Ashraf³, Aqsa Shoukat⁴, Ahmad Nawaz³, Kumail Hassan³

¹Department of Anatomy, Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan.

²Department of Parasitology, Faculty of Veterinary Science, University of Agriculture, Faisalabad, Punjab, Pakistan.

³Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan.

⁴Department of Pathology, Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan

*Correspondence: Sabarashid440@gmail.com

ARTICLE INFO

ARTICLE HISTORY: CVJ-24-0104

Received 10 January 2024
Revised 15 April 2024
Accepted 05 May 2024
Published 30 June 2024
online

Keywords:

Resistance
Phytomedicine
Poultry
Herbal
plants

ABSTRACT

With the advancing population, poultry has attained a valuable food and economic position. It has become a necessary business, from scavenging backyard poultry to the \$7.87 billion net worth of JBS poultry. Any kind of minor disturbance may cause a high strain on poultry production and parasitic infections are impacting the poultry industry in terms of production and management. As directly concerned with the gastrointestinal tract, parasitic infections hamper the feed conversion ratio, which is the base of poultry farming. Drugs used to treat these parasitic infections are losing their sensitivity because of various factors, including unchecked and repetitive use. Veterinary ethnopharmacology is adapting towards the use of botanicals, essential oils, and natural remedies to treat these parasitic infections. Recent studies have proved the efficacy of these products. This review aims to provide a brief comparison of these herbal products with anthelmintics, their efficacy, and their future in ethnopharmacology.

To Cite This Article: Rashid S, F Hafeez, R Ashraf, A Shoukat, A Nawaz and K Hassan, 2024. Phytomedicine efficacy and prospects in poultry; a new insight to old anthelmintic resistance. Continental Vet J, 4(1):62-75.

Introduction

Food animals have always been a source of greater attention, as they are concerned with public health, consumer demand, and the metabolic kinetics of consumers. With the elevating population, an increase in demand for food is natural. To meet this increasing food demand, there has been an increase in the animal population especially in tropic and sub-tropic areas (Osti et al. 2017). Poultry has been used as a source of food since the evolution of humankind, as it is an at-hand source of protein available (Khan 2018). To fulfill the required poultry production, especially broiler production is expected to meet the critical storage of animal protein need (Hatab et al., 2019; Kpomasse et al. 2021).

Poultry is the second largest industry in Pakistan and 11th largest industry in the world (Rehan et al. 2019). According to FAO statistics (2012-2022), poultry egg and meat demand is the highest among all the meat types (Desier et al. 2018). Various production systems have been introduced in poultry, i.e. cage system systems, free range, backyard poultry, etc. (Karcher and Mench, 2018). In all these systems, the process of disinfection, medication, vaccination, environmental control, manpower involvement, the progress of the disease, worm infestation, and all these factors are variable. The course of the disease is quite low in production systems, where an all-in-all-out system is applied (Zhou et al. 2020) and the feeding is controlled. When dealing with poultry, worm infestation is a major issue causing economic, commercial, and

health losses. This worm infestation is observed highly in organic poultry and backyard poultry. It is because of their higher exposure to the environment. Economically, parasitic infections are of high importance (Blake et al. 2020). Moreover, the chances of parasitic infections in deep litter poultry systems and other production systems have also been reported (Singh et al. 2021). As parasites directly interact with the gastrointestinal tract of poultry, serious impacts are posed on the feed conversion ratio (FCR) and performance, which in turn impacts the economics of poultry farming (Ahmad et al. 2022). Effects on Feed conversion ratio (FCR) and performance mean production losses and production losses refer to less egg production in layers and poor weight gain in broilers. The probability of helminth infection varies from system to system (Sharma et al. 2019). However, the occurrence of parasitic infection is quite high in organic poultry and free-range systems (McMullin 2022). Because of free access to the environment and open fields, this type of poultry is more likely to have worm infestation. According to some reports, Even the eggs of *Taenia solium* have also been reported in the eggs of organic poultry (Dixon et al. 2020). The probability of *Ascaridia galli* is the highest in backyard poultry and free-range systems (Tomza-Marciniak et al. 2014). Anthelmintic drugs are widely used and prepared in veterinary medicine because of this huge concern for parasitic infection (Castagna et al. 2021). But their consequences have also become a source of interest. Poultry parasites have become resistant to these anthelmintic drugs. This means parasites do not show any sensitivity or response to these drugs when used (Sharma et al. 2019). Many factors are involved in the development of anthelmintic drug resistance. The genes of nematodes can change, with less or incorrect concentration of drug dose, specific causes, experimental use of drugs, efficacy of drug, etc. Some studies have also revealed the toxic impacts of anthelmintics as their residues were found in poultry (Zirintunda et al. 2022). All these things demand an alternative to be used efficiently as anthelmintic. Ethnoveterinary practices have always been there in our field. Essential oils and medicinal plants are used as anthelmintics (Ramdani et al. 2023). Plants have strong antioxidant impacts that not only trigger the life status of animals but also have astringent properties (Quideau et al. 2011). Plants have secondary metabolites like polyphenols, tannins, and alkaloids that exhibit medicinal properties of plants. Many plants have been successfully used as anthelmintic without even knowing the specific metabolite or active ingredient responsible for their anthelmintic action (Oliveira et al. 2021). This review aims to review the anthelmintic activity of plants used as anthelmintic, their mode of action, pros and cons, and efficacy against parasites.

Helminthes infections in poultry

While raising poultry, external or internal parasites pose serious impacts on the quality as well as quantity of production (Gržinić et al. 2023). Internal parasites of helminths predilection sites are mostly in the gastrointestinal tract (Jilo et al. 2022). While raising poultry, production cost and FCR are two major points to be considered for successful and efficient production (Muhammad et al. 2023). Major economic losses are caused by *Ascaridia galli*, *Heterakis gallinarum*, and *Eimeria* species of parasites. The most important infection of poultry is coccidiosis which is caused by *Eimeria* species having a mortality rate of 50%. Moreover, coccidiosis also promotes certain other infections and provides an environment for *Clostridium perferingens* (Abd El-Hack et al. 2022). Blackhead, a disease of the lower digestive tract is also caused by *Histomonas meleagridis*. Acaridiasis is caused by *Ascaridia galli* which is a nematode. It is also a disease of high economic importance. It impacts egg production loss and reduced growth rate as in the presence of *Ascaridia galli* in the intestine, nutrient absorbance is at its lowest (Abon 2021). Some important helminths of poultry are mentioned in Table 1.

Commonly used anthelmintics in poultry

While considering the mortality rates and economic losses, certain types of anthelmintic are used (Kadykalo et al. 2018). The efficacy of these anthelmintics has been analyzed *in vitro* and *in vivo*, by induced infections or natural infections (Bazh and El-Bahy 2013). The infective stages, time of anthelmintic administration, and dosage were checked against commonly present helminths (Williams et al. 2014).

The parameters for determining the potential of anthelmintics are necropsy findings, fecal and cecal egg count, worm burden, and number of larvae (Nielsen et al. 2022). The compounds for helminthiasis approved by the USA are given in Table 2.

Mechanism of action of anthelmintic

The activity of anthelmintic depends upon the specific anthelmintic used. The mode of action varies from compound to compound. The major site of target in anthelmintic is the cellular integrity and neuromuscular degeneration of the parasite (Holden-Dye and Walker 2014). Whereas the specific site of action may vary. The process by which cellular integrity is affected is explained in Fig. 1.

Macrocyclic lactones: These drugs work by binding the glutamate-gated chloride channels present in the nerve cells (Wolstenholme et al. 2016). This binding causes the channel to allow an agonistic activity towards worm muscle receptors

i.e. L-subtype nicotinic acetylcholine receptors (Weeks et al. 2018).

Monepantel: It is the amino-acetonitrile derivative, it is the agonist of mptl-1 channel of the DEG-3 family of acetylcholine receptors (Baur et al. 2015). These receptors are present in nematodes only and lead to muscle depolarization and irreversible paralysis.

Derquantel: It is like piperazine causes flaccid paralysis in nematodes. But in the case of derquantel, it happens by inhibiting 45-pS channels (Tuck et al. 2016).

Piperazine: Another anthelmintic that works by hyperpolarizing the nerve membranes, thus breaking neuromuscular links and causing flaccid paralysis (Puttachary 2012)

Table 1: Common helminths of poultry, site of predilection, mortality rate and prepatency period

Parasite	Life Cycle	Predilection Site	Pathogenicity	Mortality Rate (%)	Prepatency period (Days)
Nematodes					
<i>Amidostomum anseri</i>	Direct	Gizzard	Severe	Up to 100	14-25 days
<i>Ascaridia galli</i>	Direct	Small intestine	Moderate	40	45 days
<i>Eucoleus annulata</i>	Earthworm	Esophagus, crop	Moderate to severe	>10	3-4 weeks
<i>Eucoleus contortus</i>	None or earthworm	Mouth, esophagus, crop	Severe	31	3-4 weeks
<i>Baruscapillaria obsignata</i>	Direct	Small intestine, ceca	Severe	30-35	19-22 days
<i>Cyrnea colini</i>	Cockroach	Proventriculus	mild	<1	5-6 weeks
<i>Heterakis gallinarum</i>	Direct	Ceca	Mild, but transmit histomoniasis	10	7-12 days
<i>Heterakis isolonche</i>	Direct	Ceca	Severe	50	24-30 days
<i>Strongyloides avium</i>	Direct	Ceca	Moderate	60	10 days
<i>Trichostrongylus tenuis</i>	Direct	Ceca	Severe	44	5-10 days
Cestodes					
<i>Choanotaenia infundibulum</i>	House flies	Upper intestine	Moderate	40.67	2-4 weeks
<i>Raillietina cesticillus</i>	Beetles	Duodenum, jejunum	Mild	52.6	28 days
<i>Raillietina tetragona</i>	Ants	Lower intestine	Severe	45	13 days
<i>Davainea proglottina</i>	Snails	Duodenum	Severe	37	2-3 weeks
Trematodes					
<i>Postharmostomum commutatum</i>	Unknown	Caecum	Moderate to severe	10	14 days
<i>Zygocotyl lunata</i>	Snails	Caecum	Severe	20	60 days
<i>Hypoderaeum conideum</i>	Snails, fishes	Posterior small intestine	Mild	30	3 weeks

a great influx of chloride ions. These chloride ions accumulate at different sites with different sensitivities. As a result, the effect of macrocyclic compounds varies. The pharynx is paralyzed, so the mobility of the digestive tract is lost. And even if the paralysis recovers due to lower sensitivity of chloride ions, the feeding intake not regained due

to pharynx paralysis that contributes to helminths death.

Emodepside: These drugs act by the stimulation of latrophilin-like receptors, which are present at the neuromuscular junction synapse (Holden-Dye et al. 2012). These receptors transduce signals to Gq-

proteins and phospholipase C, due to which internal calcium and diacylglycerol levels are increased and calcium-activated potassium channel slo-1 is targeted. Binding to the latrophilin receptor and the slo-1 ion channel leads to inhibition of pharyngeal pumping, paralysis and death.

Development of Resistance

Resistance against anthelmintic can be described in the terms that helminths lose sensitivity to a specific chemical used against it (Fissiha and Kinde 2021). Resistance is one of the most common issues faced while dealing with the helminths in poultry. Resistance develops due to various causes. It is due to incorrect dosage and administration (Kaye and Pogue 2015). Most of the helminths are shed in feces of poultry. In backyard poultry, it is a common practice for poultry to peck other feces. This repeated exposure to helminths and anthelmintic causes the resistance to develop (Shalaby 2013). Low potent drugs, and incorrect knowledge of the specific helminths is also contributing towards resistance. certain helminths are resistant to certain anthelmintics, e.g. fenbendazole is resistant against *Ascaridia dissimilis*, and *Heterakis gallinarum*, *Histomonas meleagridis* (Patel et al. 2018). Another important contributing factor is the unavailability of any diagnostic test or instruments for checking resistant. Sometimes resistance develops because of mutations in bgenes(Raza genes (Raza et al. 2016). Moreover, as with macrolide compounds, the sensitivity of chlorine receptors varies, and the concentration of chlorine ion influx causes varying concentrations at various sites. This unknown accumulation may also lead to resistance.

Phytomedicine

The use of plants and plant products in medicine for the treatment of various infections either

parasitic, bacterial, or viral is phytomedicine. It is one of the most explored research areas nowadays because of its high efficacy, proficiency, and availability. Herbal products are believed to cause very little harmful effects on the health of animals. The residues of botanicals in the meat and other products of food animals are very uncommon. Botanicals are now being used in various diseases widely, including parasitic diseases. Many of the parasites of poultry have become resistant to previously used anthelmintics because of multiple unattended causes. Concerning the economic importance of poultry, efforts are quite noticeable in dealing with the anthelmintic resistance (Wolstenholme 2011). As anthelmintic resistance can pose serious impacts on commercial meat production and meat quality. Several medicinal plants have been tested against resistant helminths infections *in vitro* and *in vivo* (Romero-Benavides et al. 2017). Beyond less cost, medicinal plants are more effective, highly potent, have no toxicity, and have fewer residues in meat. This may pose long term effects on the economy of poultry farmers specifically in developing countries (Conan et al. 2012).

Phytomedicine and Helminthiasis

Anthelmintics have characteristics like cost effectiveness, single-dose cure and spectrum broad-spectrum activity (Panic et al. 2014). Botanicals used in helminthiasis have active ingredients such as Polyphenols, saponins, tannins, alkaloids, isoquinoline, maloinic acid, Proanthocyanidin, gallic acid, ellagic acid, flavonoids (Jamil et al. 2022). These compounds have their specific mode of action against specific anthelmintics. Mostly ant anthelmintics work by disrupting the neuromuscular activity of larvae, as anthelmintic resistance passes in genes so disrupting the worm infestation at a very early stage is required to overcome helminthiasis (Shalaby 2013).




Fig. 1: Mode of action of anthelmintics in cellular integrity impairment

Saponins-based herbal anthelmintics

Saponins include steroids, ginsenosides, allium saponins, triterpenoids, and glycoalkaloid saponins (Soetan et al. 2014). There are 11 classes of saponins usually present on plants. Plant saponins which are used as anthelmintics are found in Aloeaceae, Amaranthaceae, Zygophyllaceae, Rosaceae, Fabaceae, Apiaceae, and Verbenaceae (Oakenfull and Sidhu 2023.) Saponins are used in fungal, parasitic, and bacterial infections. Saponins that have anthelmintic properties listed in Table 3 work by disrupting the mitochondrial action resulting in low cellular energy (Faixová et al. 2021), and inability to move, which causes the death of helminths (Fig. 2). Some saponins work by increasing the permeability of the helminth cell membrane (Singla and Kaur 2021), allowing the passage of certain undesirable compounds which cause helminth to die (Fig. 2).




Fig. 2: Mechanism of action of saponins based saponins

Terpenoids

Terpenoids include monoterpenes, diterpenes, triterpenes, sesterpenes and sesquiterpenes (Perveen 2021). It is made up of repeating units of carbon. The major source of terpenes is from the leaves of eucalyptus tree. Digoxin, lupeol, ursolic acid, oleanolic, atrastyloside, cicutoxin, atractyloside, and betulinic acid, are the different types of terpenoids (Surbhi et al. 2023). All these

types interfere with helminths infestation in different ways, by muscle contraction (varsha et al. 2017), increasing calcium ion absorption, blocks oxidative phosphorylation (Mbaveng et al. 2014), decreasing antioxidant enzymes, inhibiting ATP synthesis (Baliga et al. 2019), increasing cytochrome C by mitochondria (Dev et al. 2017) and lowering glucose levels of helminths (Fig. 3). The most common terpenoids listed in table 4 used for their anthelmintic properties are those from camphor and methanol (Mukherjee et al. 2016). Terpenoids interfere with the neurotransmission, leading to paralysis of helminth and ultimate death (Mukherjee et al. 2016). Terpenes are known to be effective against albendazole-resistant helminths (Mirza et al. 2020).

Flavonoids-based anthelmintics

Flavonoids are polyphenolic compounds that have enzymatic activity such as phosphodiesterase and Ca²⁺ ATPase (Al-sanafi 2020). Flavonoids are known for their synergistic action with botanical anthelmintics such as tannins and other anthelmintics like praziquantel (lateef et al. 2013). Flavonoids are known for their anticancer, inflammatory, and antioxidant effects (Liu et al. 2020). Flavonoids are known for reducing the EPG, juvenile and adult worms' survival rate.

Tannins and alkaloids-based anthelmintics

Tannins are polyphenols with many medicinal properties (sahakyan et al. 2020). Tannins also work as anthelmintics and paralyze the worm's causing death. Condensed tannins are usually known as anthelmintics. They need to be administered with care as they are antinutritional. Tannins are environmentally stable and degrade at very high temperatures (Jindal and Nikhanj 2023). Alkaloids are used as cardio protectors, anesthetics, and anti-inflammatory agents. Alkaloids are the neurotoxin anthelmintics, mostly work by breaking the cell walls, protein, and lipid content of worms (Molan 2014). Usually, the most common alkaloid anthelmintic is steroidal glycoside solamargine from plants. Glucosinolates interact with the DNA integrity of cells, acacia oxyphylla cause flaccid paralysis (Ade-Ademilua and okpoma 2018).




Fig. 3: Possible mechanisms of actions of terpenoids on helminths

Efficacy of botanical anthelmintic

Demand for anthelmintic alternatives is increasing rapidly due to certain causes like resistance, high-cost products, and less availability (Sepulveda-Crespo et al. 2020). At least studies have been conducted on the pharmacokinetics and pharmacodynamics of plants. Their efficacy can only be measured by conducting studies on the pharmacological impact of these plants. But their availability, less cost, need for alternative is making them highly susceptible in developing countries,

specifically Asian countries (Nelson et al. 2019). The efficacy of anthelmintic drugs and their chemotherapy has always been a problem, due to several species, genetic variation in helminths species, developing resistance, toxic effects, and residues in meat and eggs. So, the need for alternatives that can cover all these impacts is quite important. Unfortunately, phytomedicines have given much less importance in research, as compared to their need




Fig. 4: Mechanisms of actions of commonly used anthelmintic drugs

Future prospectives

If proper pharmacological research is carried out on the impact of botanicals use as anthelmintic. In the upcoming years botanicals will completely replace anthelmintic drugs. Using these nonresistant anthelmintics can save the economic losses occurring from nematode infections yearly. Coccidiosis was able to cause 3 billion US dollar losses every year. Similarly, ascardiasis in poultry causes 10-40% loss in egg production. These all can be efficiently controlled by using botanicals. Moreover, residues in the meat or poultry after using anthelmintic drugs are also a major loss in broilers. Being considering poultry as a huge industry, its economics is taken seriously and that is improved by using quality nonresistant botanical anthelmintic.

Funding

No funds were provided by any national or international funding agencies.

Ethical statement

No Ethical permissions were required for this article.

Availability of data and material

The data can be obtained from the corresponding author on a reasonable request.

Acknowledgement

We acknowledge the technical support of the Department of Zoology, The Islamia University, Bahawalpur, for publication of this article.

Consent to participate

All the authors gave their consent for equal participation.

Consent for publication

All the authors gave their consent for publication.

Competing Interest

The authors declare that they have no relevant financial or non-financial interests to disclose.

Author Contribution

MR, AA, and BES wrote the manuscript NA, YM, and GM managed references, tables, and figure, TS and MF revised the article.

Table 4: Terpenoids used as anthelmintics in plants, their composition, MOA and Method of extraction

Medicinal plant	Common name	Part of plant used	Active compound	MOA	Composition	Method of preparation	References
<i>Artemisia afra</i>	African wormwood	Leaves	Flavonoids	Block the neuromuscular transmission in a parasite by depolarizing nerve impulse conduction	Terpenoids, essential oils, coumarins, caffeoylquinic acid	Hexane extraction	(Symeonidou et al. 2018)
<i>Dryopteris filix-mas</i> Schott	Male fern	Leaves, stems	Quercetin	Multi target mechanism.	Terpenoids	Maceration with aqueous ethanol.	(Ramlal et al. 2023)
<i>Momordica charantia</i> L	Bitter melon	Fruit	Charantin, vicine.	Slow down the process of gluconeogenesis (Ramlal et al. 2023).	Sterols, terpenoids.	Filtration and evaporation in reduced pressure.	(Liu et al. 2020)
<i>Mentha longifolia</i>	Wild mint	Leaves	Pulegone	Inhibit the NF-kB signaling pathway	Menthol, cineole, menthone	Hydro distillation technique using Soxhlet apparatus.	(Lateef et al. 2023)
<i>Curcuma longa</i>	Turmeric rhizome	Rhizome extract	Curcumin	Interfere with Nf-kB (Ramlal et al. 2023)	Curcumin, demethoxycurcumin, bisdemethoxycurcumin	Hexane based extraction, hydro distillation.	(Manjusa and Pradeep 2022)
<i>Cucurbita moschata</i> Duchesne	Pumpkin	Seeds	Cucurbitin	Disrupt the neurotransmission and disruption of helminths egg	Carbohydrates, proteins, lipids, fiber	Organic solvent assisted extraction	(Lateef et al. 2013)

Table 2. Commonly used anthelmintic in poultry, target specie, mode of action, and percentage efficacy

Anthelmintic	Commonly infected species	Dosage	Withdrawal Time	Percentage efficacy (%)	Mechanism of Action
Piperazine compounds	<i>Ascaridia</i> spp.	Chickens: Single dose, 50 mg/bird (< 6 weeks old), 100 mg/bird (≥6 weeks old), in the feed at 0.2%–0.4% or in the drinking water at 0.1%–0.2% Turkeys: Single dose 100 mg/bird (< 12 weeks old) or 200 mg/bird (≥12 weeks old) Because only the piperazine moiety is efficacious, doses should be calculated based on mg of active piperazine/bird. Treatment can be repeated after 14 days.	14 days (meat); not approved in the USA for birds producing eggs for human consumption.	99-10	It binds with gamma butyric acid receptor subunit beta-3 which causes hyperpolarization of nerve endings and paralyzes the worm by this live worm s excluded from the body.
Fenbendazole	<i>Ascaridia</i> spp, <i>Heterakis gallinarum</i>	Growing turkeys: 14.5 g/ton of feed (16 ppm), fed continually as the sole ration for 6 days.	None for turkeys Not approved for other poultry.	More than 80	It binds with the tubulin protein of parasites thus inhibit the production of microtubules and poses negative impact on the transport system of parasite.
Hygromycin B	<i>Ascaridia</i> spp, <i>Heterakis gallinarum</i> , <i>Capillaria</i> spp.	8–12 g/ton in feed	3 days for meat	0-100	Inhibit Protein synthesis, growth rate, affect 30S ribosomes
Coumaphos	<i>Capillaria</i> spp.	Replacement pullets: 0.004% in feed for 10–14 days Layers: 0.003% in feed for 14. In birds maintained on contaminated litter or exposed to infected birds, the medication regimen should be repeated using the same dosages as the first. There should be at least 3 weeks between the first and second course of treatment.	Unknown	88.6	Immobilize nervous system of parasite
Thiabendazole	<i>Syngamus trachea</i>	Pheasants: 0.05% in feed for 14 days	21 days for meat	0-9	Inhibit the helminth specific enzyme fumarate reductase flavoprotein subunit

Table 3: Saponins used as anthelmintic in plants, their composition, MOA and Method of extraction

Medicinal plant	Common name	Part of the plant used	Active compound	MOA	Composition	Method of preparation	References
<i>Cyamopsis tetragonoloba</i> Taub	Guar bean	Flowers oil	Saponins	Saponins lyse the oocyst of helminths.	Ethyl acetate and methanolic acid	Grinding of endosperm portion	(Abo-EL-Sooud 2018)
<i>Helichrysum splendidum</i> less	Mulberry pines	Leaves, flowers	saponins	Inhibit the gene expression of NO (Lateef et al. 2013)	Phellandrene, cadinene, germacerene, cineole.	Microwave extraction technique	(Veerakumari 2015)
<i>Sena occidentalis</i>	Coffee sienna	Leaves	Saponosides	Hyperpolarizing of the nerve membrane, which leads to flaccid paralysis in helminths (Symeonidou et al. 2018)	Galactomannan, polysaccharides, glycosides	Ethanol-based leaf extraction	(Saeed and Alkheraije 2023)
<i>Tehrosia Vogelli</i> Hook.f	Fish poison bean	Leaves, stems	Deguelin	Inhibit larval development and motility.	saponins, alkaloids, phenols.	Hexane and ethanol-based extraction.	(Veerakumari 2015)
<i>Momordica corantia</i> L	Bitter melon	Fruit	Charantin, vicine.	Slow down the process of gluconeogenesis ((Ramlal et al. 2023).	Sterols, terpenoids, saponins	Filtration and evaporation in reduced pressure.	(Liu et al. 2020)
<i>Morinda lucida</i>	Brimstone tree	Leaf and bark extract	Alkaloids	Reduce nitric oxide and E-2 levels	Saponins, polyphenols, reducing sugars,	Methanol based maceration	(Saeed and Alkheraije 2023)
<i>Echinacea purpurea</i> Moench	Coneflower	Root, leaves and flowers	Chloric Acid	Activate phagocytosis and increased respiratory activity	Polysaccharides, lipoproteins, saponins, betaine	Supercritical carbon dioxide extraction	(Abo-EL-Sooud 2018)
<i>Carica papaya</i> L	Paw paw	Seeds	alkaloids, saponins	Activation of caspase-3/7 pathway, increase peripheral glucose, NF-kB attenuation	BITC, carbohydrates, proteins, proteolytic enzymes	Boiling and shaking	(Lateef et al. 2013)

Table 5: Flavonoids used as anthelmintic in plants, their composition, MOA, and Method of extraction

Medicinal plant	Common name	Part of plant used	Active compound	MOA	Composition	Method of preparation	References
<i>S. flavescens</i>	Shrubby sophora	Root extract	Sophorae Radix	Induce apoptosis and autophagy by MCF-7 cells (Kostadinovi, et al. 2015)	Alkaloids and flavonoids	Ethanol purification followed by ion exchange resin absorption	(Veerakumar 2015)
<i>Allium sativum</i>	garlic	cloves	Organosulfur compounds	Modulate the expression of Nf-kB transcription factors	Flavonoids, alliin, ajoenes, allicin	Boiling with ethanol extracts	(Liu et al. 2020)
<i>Moringa oleifera</i>	Drumstick tree	Seed extract	Flavanol Rutin	Inhibit the formation and mobility of larvae.	Carbohydrate, protein, fat, fiber	Superficial extraction with CO ₂ and ethanol.	(Lateef et al. 2013)
<i>Punica Granatum L.</i>	Pomegranate	peels	Ellagic acid	Disrupt the neurotransmission leading to paralysis of larvae	Anthocyanins, flavonoids, fructose, sucrose, organic sugars	High pressure nonpolar extraction	(Manjusa and Pradeep 2022)
<i>Flemingia vestita L</i>	Sohphlang	Root tuber	Genistein	Alters the glycolytic enzymes of cestodes (Jamil et al. 2022)	<i>Genistein, pseudobaptigenin, daidzein.</i>	Ethanol extraction	(Liu et al. 2020)
<i>Vitis vinifera</i>	Grape seed	Fruit	Proanthocyanidin	Act on the IL-8 promoter activity	Tannins, proteins, cellulose, lignin, hemicellulose	Maceration and elevated temperature extraction	(Liu et al. 2020)
<i>Camellia sinensis kuntze</i>	Green tea	leaves	polyphenols	Inhibition of intracellular enzymes, cell wall and cell membrane disruption	Tannin, Epigallocatechin-2, Gallate.	Ethanol extract	Bora and Bora (2021)

References

- Abd El-Hack ME, El-Saadony MT, Salem HM, El-Tahan AM, Soliman MM, Youssef GB, Taha AE, Soliman SM, Ahmed AE, El-Kott AF and Al Syaad KM, 2022. Alternatives to antibiotics for organic poultry production: types, modes of action and impacts on bird's health and production. *Poultry Science*, 101(4): 101696.
- Abd El-Hack ME, El-Saadony MT, Elbestawy AR, Nahed A, Saad AM, Salem HM, El-Tahan AM, Khafaga AF, Taha AE, AbuQamar SF and El-Tarabily KA, 2022. Necrotic enteritis in broiler chickens: disease characteristics and prevention using organic antibiotic alternatives—a comprehensive review. *Poultry Science* 101(2): 101590.
- Abon AC, 2021. Anthelmintic efficacy of tuba (*Croton tiglium* L.) seeds on the gastrointestinal parasites of native chickens (*Gallus domesticus*). *Plant Science Today* 8(4): 749-753.
- Abo-EL-Sooud K, 2018. Ethnoveterinary perspectives and promising future. *International Journal of Veterinary Science and Medicine* 6(1): 1-7.
- Ade-Ademilua OE and Okpoma MO, 2018. *Gossypium Hirsutum* L. And *Gossypium barbadense* L; Differences in Phytochemical Contents, Antioxidant and Antimicrobial Properties. *Life Sciences Journal*. 20 (1), 77. doi:10.4314/ijls.v20i1.8
- Ahmad R, Yu YH, Hsiao FSH, Su CH, Liu HC, Tobin I, Zhang G and Cheng YH, 2022. Influence of heat stress on poultry growth performance, intestinal inflammation, and immune function and potential mitigation by probiotics. *Animals* 12(17): 2297.
- Al-Snafi AE, 2020. Phenolics and flavonoids contents of medicinal plants, as natural ingredients for many therapeutic purposes-A review. *IOSR Journal of Pharmacology* 10(7): 42-81.
- Blake DP, Knox J, Dehaeck B, Huntington B, Rathinam T, Ravipati V, Ayoade S, Gilbert W, Adebambo AO, Jatau ID and Raman M, 2020. Re-calculating the cost of coccidiosis in chickens. *Veterinary Research* 51: 1-14.
- Bora P and Bora LC, 2021. Microbial antagonists and botanicals mediated disease management in tea, *Camellia sinensis* (L.) O. Kuntze: an overview. *Crop Protection* 148: 105711.
- Baliga MS, Shivashankara AR, Venkatesh S, Bhat HP, Palatty PL and Bhandari G, 2019. Phytochemicals in the Prevention of Ethanol-Induced Hepatotoxicity: A Revisit. *Dietary Interventions Liver Disorders. Foods Nutrients, Dietary Supplements* 2019: 79-89.
- Baur R, Beech R, Sigel E and Rufener L, 2015. Monepantel irreversibly binds to and opens *Haemonchus contortus* MPTL-1 and *Caenorhabditis elegans* ACR-20 receptors. *Molecular Pharmacology* 87(1): 96-102.
- Castagna F, Piras C, Palma E, Musolino V, Lupia C, Bosco A, Rinaldi L, Cringoli G, Musella V and Britti D, 2021. Green veterinary pharmacology applied to parasite control: Evaluation of *Punica granatum*, *Artemisia campestris*, *Salix caprea* aqueous macerates against gastrointestinal nematodes of sheep. *Veterinary Sciences* 8(10): 237.
- Conan A, Goutard FL, Sorn S and Vong S, 2012. Biosecurity measures for backyard poultry in developing countries: a systematic review. *BMC Veterinary Research* 8: 1-10.
- Dev K, Ramakrishna E and Maurya R, 2017. Glucose Transporter 4 Translocation Activators from Nature. *Discovery and Development of Antidiabetic Agents from Natural Products. Drug Discovery* 2017: 113-145.
- Desiere S, Hung Y, Verbeke W and D'Haese M, 2018. Assessing current and future meat and fish consumption in Sub-Sahara Africa: Learnings from FAO Food Balance Sheets and LSMS household survey data. *Global Food Security* 16: 116-126.
- Dixon MA, Winskill P, Harrison WE, Whittaker C, Schmidt V, Sarti E, Bawm S, Dione MM, Thomas LF, Walker M and Basanez MG, 2020. Force-of-infection of *Taenia Solium* porcine cysticercosis: a modelling analysis to assess global incidence and prevalence trends. *Scientific Reports* 10(1): 17637.
- Faixová D, Hřčková G, Kubašková TM and Mudroňová D, 2021. Antiparasitic effects of selected isoflavones on flatworms. *Helminthologia* 58(1): 1-16.
- Fissiha W and Kinde MZ, 2021. Anthelmintic resistance and its mechanism: A Review. *Infection and Drug Resistance* 5403-5410.
- Gržinić G, Piotrowicz-Cieślak A, Klimkowicz-Pawlas A, Górny RL, Ławniczek-Wałczyk A, Piechowicz L, Olkowska E, Potrykus M, Tankiewicz M, Krupka M and Siebielec G, 2023. Intensive poultry farming: A review of the impact on the environment and human health. *Science of The Total Environment* 858: 160014.
- Holden-Dye L, Crisford A, Welz C, von Samson-Himmelstjerna G, Walker RJ and O'Connor V, 2012. Worms take to the slo lane: a perspective on the mode of action of emodepside. *Invertebrate Neuroscience* 12: 29-36.
- Holden-Dye L and Walker R, 2014. Anthelmintic drugs and nematocides: studies in *Caenorhabditis elegans*. *WormBook: The Online Review of C. Elegans Biology*; pp: 1-29.

- Jamil M, Aleem MT, Shaukat A, Khan A, Mohsin M, Rehman TU, Abbas RZ, Saleemi MK, Khatoon A, Babar W and Yan R, 2022. Medicinal plants as an alternative to control poultry parasitic diseases. *Life* 12(3): 449.
- Jindal P and Nikhanj P, 2023. A review on processing technologies for value added millet products. *Journal of Food Process Engineering* 46(10): 144-19.
- Jilo SA, Abadula TA, Abadura SZ, Gobana RH, Hasan LA and Nair SP, 2022. Review on epidemiology, pathogenesis, treatment, control and prevention of gastrointestinal parasite of poultry. *International Journal of Veterinary Sciences and Animal Husbandry* 7(5): 26-34
- Joshi B, Panda SK, Jouneghani RS, Liu M, Parajuli N, Leyssen P, Neyts J and Luyten W, 2020. Antibacterial, antifungal, antiviral, and anthelmintic activities of medicinal plants of Nepal selected based on ethnobotanical evidence. *Evidence-Based Complementary and Alternative Medicine* 2020(1); pp: 1043471
- Karcher DM and Mench JA, 2018. Overview of commercial poultry production systems and their main welfare challenges. In *Advances in poultry welfare*. Woodhead Publishing; pp: 3-25
- Kaye KS and Pogue JM, 2015. Infections caused by resistant gram-negative bacteria: epidemiology and management. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* 35(10): 949-962.
- Kumarasingha R, Palombo EA, Bhave M, Yeo TC, Lim DSL, Tu CL, Shaw JM and Boag PR, 2014. Enhancing a search for traditional medicinal plants with anthelmintic action by using wild type and stress reporter *Caenorhabditis elegans* strains as screening tools. *International Journal for Parasitology* 44(5): 291-298.
- Khan SH, 2018. Recent advances in role of insects as alternative protein source in poultry nutrition. *Journal of Applied Animal Research* 46(1): 1144-1157.
- Kostadinović L, Puvača N, Popović S and Lević J, 2015. Botanical supplements as anti-coccidial alternatives in poultry nutrition. *World's Poultry Science Journal* 71(1): 27-36.
- Lateef M, Iqbal Z, Sajid MS, Abbas RZ, Sindhu ZUD, Akhtar M, Khan MN, Awais MM, Iqbal A and Ain QU, 2013. An account of botanical anthelmintics and methods used for their evaluation. *Review Veterinary Animal Science* 1(1): 6-14.
- Liu M, Panda SK and Luyten W, 2020. Plant-based natural products for the discovery and development of novel anthelmintics against nematodes. *Biomolecules* 10(3): 426.
- Manjusa A and Pradeep K, 2022. Herbal anthelmintic agents: a narrative review. *Journal of Traditional Chinese Medicine* 42(4): 641.
- Mukherjee N, Mukherjee S, Saini P, Roy P and Sinha Babu PS, 2016. Phenolics and Terpenoids; the Promising New Search for Anthelmintics: A Critical Review. *Mini-Reviews Medicinal Chemistry* 16 (17): 1415-1441.
- Mbaveng AT, Hamm R and Kuete V, 2014. Harmful and Protective Effects of Terpenoids from African Medicinal Plants. *Toxicology Survey of African Medicinal Plants*, Elsevier; pp: 557-576.
- Mirza Z, Soto ER, Hu Y, Nguyen TT, Koch D and Aroian RV, 2020. Anthelmintic Activity of Yeast Particle-Encapsulated Terpenes. *Molecules* 25 (13): 2958.
- McMullin P, 2022. Infectious diseases in free-range compared to conventional poultry production. *Avian Pathology*, 51(5): 424-434.
- Molan AL, 2014. Effect of purified condensed tannins from pine bark on larval motility, egg hatching and larval development of *Teladorsagia circumcincta* and *Trichostrongylus colubriformis* (Nematoda: Trichostrongylidae). *Folia Parasitologica* 61(4): 371.
- Muhammad T, Ullah F, Jelani G and Atif M, 2023. A Comparative Analysis of the Growth Rate, Body Weight Gain, and Feed Conversion Ratio of Broilers Raised Under Ad Libitum and Restricted Feeding Programs. *Indus Journal of Agriculture and Biology* 2(1): 1-9.
- Nelson GC, Rosegrant MW, Koo J, Robertson R, Sulser T, Zhu T, Ringler C, Msangi S, Palazzo A, Batka M and Magalhaes M, 2009. Climate change: Impact on agriculture and costs of adaptation. *International Food Policy Research Institute*.
- Osti R, Bhattarai D and Zhou D, 2017. Climatic variation: effects on stress levels, feed intake, and bodyweight of broilers. *Brazilian Journal of Poultry Science* 19: 489-496.
- Oliveira M, Lima CS, Ketavong S, Llorent-Martínez EJ, Hoste H and Custódio L, 2021. Disclosing the bioactive metabolites involved in the in vitro anthelmintic effects of salt-tolerant plants through a combined approach using PVPP and HPLC-ESI-MSn. *Scientific Reports* 11(1): 24303.
- Oakenfull D and Sidhu GS, 2023. Saponins. *Toxicants of Plant Origin*, Taylor and Francis; pp: 97-142.
- Panic G, Duthaler U, Speich B and Keiser J, 2014. Repurposing drugs for the treatment and control of helminth infections. *International Journal for Parasitology: Drugs and Drug Resistance* 4(3): 185-200.

- Patel T, Marmulak T, Gehring R, Pitesky M, Clapham MO and Tell LA, 2018. Drug residues in poultry meat: A literature review of commonly used veterinary antibacterials and anthelmintics used in poultry. *Journal of Veterinary Pharmacology and Therapeutics* 41(6): 761-789.
- Puttachary S, 2012. Electrophysiological studies to counter anthelmintic resistance: studies in somatic muscle and pharynx of *A. suum*, PhD Thesis, Iowa State University.
- Quideau S, Deffieux D, Douat-Casassus C and Pouységu L, 2011. Plant polyphenols: chemical properties, biological activities, and synthesis. *Angewandte Chemie International Edition* 50(3): 586-621.
- Rehan M, Aslam A, Khan MR, Abid M, Hussain S, Amber J, Anjum A and Hussain A, 2019. Potential economic impact of Newcastle disease virus isolated from wild birds on commercial poultry industry of Pakistan: A review. *Hosts Viruses* 6(1):1-15.
- Raza A, Muhammad F, Bashir S, Aslam B, Anwar MI and Naseer MU, 2016. In-vitro and in-vivo anthelmintic potential of different medicinal plants against *Ascaridia galli* infection in poultry birds. *World's Poultry Science Journal* 72(1):115-124.
- Ramlal A, Nautiyal A, Kumar J, Mishra V, Sogan N and Singab ANB, 2023. Botanicals against some important nematodal diseases: Ascariasis and hookworm infections. *Saudi Journal of Biological Sciences* 103814.
- Romero-Benavides JC, Ruano AL, Silva-Rivas R, Castillo-Veintimilla P, Vivanco-Jaramillo S and Bailon-Moscoso N, 2017. Medicinal plants used as anthelmintics: Ethnomedical, pharmacological, and phytochemical studies. *European Journal of Medicinal Chemistry* 129: 209-217.
- Ramdani D, Yuniarti E, Jayanegara A and Chaudhry AS, 2023. Roles of Essential Oils, Polyphenols, and Saponins of Medicinal Plants as Natural Additives and Anthelmintics in Ruminant Diets: A Systematic Review. *Animals* 13(4): 767.
- Sharma N, Hunt PW, Hine BC and Ruhnke I, 2019. The impacts of *Ascaridia galli* on performance, health, and immune responses of laying hens: new insights into an old problem. *Poultry Science* 98(12): 6517-6526.
- Shalaby HA 2013. Anthelmintics resistance; how to overcome it? *Iranian Journal of Parasitology* 8(1): 18.
- Sahakyan N, Bartoszek A, Jacob C, Petrosyan M and Trchounian A, 2020. Bioavailability of tannins and other oligomeric polyphenols: A still to be studied phenomenon. *Current Pharmacology Reports* 6: 131-136.
- Surbhi, Kumar A, Singh S, Kumari P and Rasane P, 2023. Eucalyptus: phytochemical composition, extraction methods and food and medicinal applications. *Advances in Traditional Medicine* 23(2): 369-380.
- Singla S and Kaur S, 2021. Anthelmintic lead compounds and their targets for drug development. *Journal of Ayurvedic and Herbal Medicine* 7(4): 265-275.
- Soetan KO, Ajibade TO and Akinrinde AS, 2014. Saponins; a ubiquitous phytochemical: a review of its biochemical, physiological and pharmacological effects. *Recent Prognosis of Medicinal Plants* 43:1-24.
- Shalaby HA, 2013. Anthelmintics resistance; how to overcome it? *Iranian Journal of Parasitology* 8(1):18.
- Sharma N, Hunt PW, Hine BC and Ruhnke I, 2019. The impacts of *Ascaridia galli* on performance, health, and immune responses of laying hens: new insights into an old problem. *Poultry Science*, 98(12): 6517-6526.
- Sohaib M and Jamil F, 2017. An insight of meat industry in Pakistan with special reference to halal meat: a comprehensive review. *Korean Journal for Food Science of Animal Resources* 37(3): 329.
- Sepúlveda-Crespo D, Reguera RM, Rojo-Vázquez F, Balaña-Fouce R and Martínez-Valladares M, 2020. Drug discovery technologies: *Caenorhabditis elegans* as a model for anthelmintic therapeutics. *Medicinal Research Reviews* 40(5):1715-1753.
- Singh M, Kaur P, Singla LD, Kashyap N and Bal MS, 2021. Assessment of risk factors associated with prevalence of gastrointestinal parasites in poultry of central plain zone of Punjab, India. *Veterinary World* 14(4): 972
- Strydom T, Lavan RP, Torres S and Heaney K, 2023. The Economic Impact of Parasitism from Nematodes, Trematodes and Ticks on Beef Cattle Production. *Animals* 13(10): 1599.
- Saeed Z and Alkherajje KA, 2023. Botanicals: A promising approach for controlling cecal coccidiosis in poultry. *Frontiers in Veterinary Science* 10: 1157633.
- Symeonidou I, Bonos E, Moustakidis K, Florou-Paneri P, Christaki E and Papazahariadou M, 2018. Botanicals: a natural approach to control ascariidiosis in poultry. *Journal of the Hellenic Veterinary Medical Society* 69(1): 711-722.
- Shah MA, Sherwani SK, Sualeh M, Kanwal S, Khan HN and Kazmi SU, 2014. In vitro anthelmintic and antiurolithic assessment of *Berberis lycium* root bark. *Journal of Pharmacognosy and Phytochemistry* 3(2): 205-210.
- Tuck S, Furey A and Danaher M, 2016. Analysis of Anthelmintic and Anticoccidial Drug Residues in Animal-Derived Foods. *Chemical Analysis of Non-antimicrobial Veterinary Drug Residues in Food* 245-309.

- Tomza-Marciniak A, Pilarczyk B, Tobianska B and Tarasewicz N, 2014. Gastrointestinal parasites of free-range chickens. *Annals of Parasitology* 60(4).
- Varsha K, Sharma A, Kaur A, Madan J, Pandey RS and Jain UK, 2017. Natural Plant-Derived Anticancer Drugs Nanotherapeutics: a Review on Preclinical to Clinical success. *Nanostructures Cancer Therapy*; pp: 775-809.
- Veerakumari L, 2015. Botanical anthelmintics. *Asian Journal of Science and Technology* 6:1881-94.
- Weeks JC, Robinson KJ, Lockery SR and Roberts WM, 2018. Anthelmintic drug actions in resistant and susceptible *C. elegans* revealed by electrophysiological recordings in a multichannel microfluidic device. *International Journal for Parasitology: Drugs and Drug Resistance* 8(3): 607-628.
- Wolstenholme AJ, Maclean MJ, Coates R, McCoy CJ and Reaves BJ. 2016. How do the macrocyclic lactones kill filarial nematode larvae?. *Invertebrate Neuroscience* 16:1-9.
- Wolstenholme AJ, 2011. Ion channels and receptor as targets for the control of parasitic nematodes. *International Journal for Parasitology: Drugs and Drug Resistance* 1(1): 2-13.
- Zhou Z, Shen B and Bi D, 2020. Management of pathogens in poultry. In *Animal Agriculture*; pp: 515-530
- Ziaei SM and Amini M, 2020. Investigation of waste in livestock and poultry industry and methods to improve feed conversion ratio in it. *ALKHAS; The Journal of Environment, Agriculture and Biological Sciences* 2(4): 5-12.
- Zirintunda G, Biryomumaisho S, Kasozi KI, Batiha GES, Kateregga J, Vudriko P, Nalule S, Olila D, Kajoba M, Matama K and Kwizera MR, 2022. Emerging anthelmintic resistance in poultry: can ethnopharmacological approaches offer a solution?. *Frontiers in Pharmacology* 12: 774896.