

Addressing Oxidative Stress and Bisphenol A (BPA) Toxicity with Antioxidant Interventions for Male and Female Reproductive Health

Adeeba Naseer^{*1,4}, Noman Waheed^{2,3}, Nusheen Ahmad⁴, Nimra Ather⁵ and Sumaira Raziq⁶

¹College of Fisheries, Southwest University, Chongqing 400715, China, ²Faculty of Fisheries, Kagoshima University, Kagoshima 890-8580, Japan, ³College of Animal Sciences, Jilin Agricultural University, Changchun, Jilin, 130118 PR, China, ⁴Department of Zoology, The Islamia University of Bahawalpur, Bahawalpur, 63100, Pakistan, ⁵Department of Zoology, Wildlife and Fisheries, University of Agriculture Faisalabad 38000, Punjab, Pakistan, ⁶Department of Zoology, Government College University Lahore 54000, Pakistan

*Correspondence: adeeba.iub22@gmail.com

ARTICLE INFO

ARTICLE HISTORY: CVJ-24-1105

Received: 05 November 2024
 Revised: 12 December 2024
 Accepted: 16 December 2024
 Published online: 19 December 2024


Key words:

Antioxidants
 Bisphenol A
 Gender toxicity
 Oxidative stress
 Reproduction

ABSTRACT

Human exposure to bisphenol A (BPA) is protracted since it is a non-persistent anthropogenic molecule that is widely used, found in numerous consumer products, and ubiquitous in the environment. This review looks at the landscape. Numerous research that have looked at the underlying molecular mechanisms of BPA toxicity over the past ten years have linked BPA-induced oxidative stress to human disease, male and female reproductive abnormalities, and other receptors in target cells due to its hormone-like property, which causes undesirable cellular reactions linked to inflammation and oxidative stress. BPA is a metabolic and endocrine disruptor that damages redox equilibrium by decreasing antioxidant enzymes and increasing oxidative mediators. This results in mitochondrial malfunction, changes to cell signaling pathways, and the activation of apoptosis. In order to comprehend how the activation of oxidative stress can be seen as the "fil rouge" of BPA's harmful modes of action with pleiotropic effects, this review will look at the current state of the BPA literature. Here, we focus on the protective effects of five classes of antioxidants vitamins and co-factors, natural products (herbals and phytochemicals), melatonin, selenium, and methyl donors (used alone or in combination) that have been found useful to counteract BPA toxicity in male and female reproductive functions.

Graphical Abstract



To Cite This Article: Naseer A, Waheed N, Ahmad N, Ather N and Raziq S 2025. Addressing Oxidative Stress and Bisphenol A (BPA) Toxicity with Antioxidant Interventions for Male and Female Reproductive Health. Continental Vet J, 5(1): 31-48. <http://dx.doi.org/10.71081/cvj/2025.033>

INTRODUCTION

Endocrine-disrupting chemicals (EDCs) are a broad category of compounds that can affect metabolic and reproductive processes as well as interfere with hormonal-signaling pathways (Sifakis et al., 2017). The most common EDC in the environment today is bisphenol A (BPA), a non-persistent anthropic chemical that has been used extensively in recent decades as a component of plastic for a variety of purposes such as food packaging, personal care, household products, and medical equipment (Meli et al., 2020). In fact, BPA has been found in food, household dust, consumer goods, aquatic ecosystems, the atmosphere, and biotic matrices. Exposure to BPA is associated with deleterious health effects for animals and humans and affects not only endocrine and reproductive organs but also immune and central nervous systems through several mechanisms, including oxidative stress (Abraham & Chakraborty, 2020). Growing evidence from research on laboratory animals shows that this non-persistent compound alters male and female reproductive function even at extremely low exposure levels. This is relevant because BPA exposure may be chronic, making it functionally equivalent to a persistent compound (Nelson et al., 2020).

The mechanisms underlying BPA toxicity may be related to its chemical properties in the body, including interactions with hormone receptors (i.e., estrogen receptors) in target cells (Cimmino et al., 2020). As a weak estrogen receptor ligand, BPA binds to the classical nuclear receptors with different affinity, for example, estrogen receptor (ER) α less than ER β (about 10,000 times lower than 17- β -estradiol), an ditspotency of action becomes higher when the estradiol level is at very low concentrations (Toporova & Balaguer, 2020). BPA also binds many other membranes and nuclear receptors, activating them at concentrations lower than those required to activate ERs (Cimmino et al., 2020). In particular, BPA binds to the membrane-bound form of estrogen receptor (mER) α and G protein-coupled receptor (GPR) 30, inducing non-genomic effects (Maniradhan & Calivarathan, 2023). GPR30 has been detected in several cell types, and it is involved in cell proliferation and apoptosis (Hernández-Silva, Villegas-Pineda, & Pereira-Suárez, 2020). Indeed, seminoma proliferation is induced by allow concentration of BPA through a GPR 30 non-genomic activation. BPA also binds to another estrogen-related receptor (ERR) γ , an orphan nuclear receptor highly expressed in the fetal brain and placenta by which BPA influences fetal tissue development (Cimmino et al., 2020).

As demonstrated by the scientific literature on BPA occurrence in human tissues and body fluids (i.e., urine, serum, plasma, saliva, breast milk, semen, follicular fluids, and adipose tissues), widespread contamination and dietary consumption have resulted in BPA exposure in the general population (Hahladakis et al., 2023). According to a recent epidemiological investigation, female reproduction may be adversely affected by elevated urine BPA concentrations in humans (Stavridis et al., 2022). Moreover, BPA exposure has been linked to

lower libido and erectile dysfunction in men, and urine BPA levels have been linked to aberrant semen parameters. When exposure happens in utero, the overall effects of BPA on male reproduction seem to be more detrimental (Pivonello et al., 2020). Animal models are crucial for evaluating the underlying toxic mechanisms seen in people, given the difficulties in both tissue sample collection and human exposure evaluation (Wetherill et al., 2007). In the first place, rats vary from humans in that they permit tissue and cell-specific molecular analysis in addition to histological evaluation of diverse tissues (Gignac & Zajenkowski, 2023).

Furthermore, rodent models offer outstanding chances for sex-specific assessments. The tissue under examination, the structural, functional, and epigenetic variations across gender and species, the duration of exposure, and the chemical dosage are all crucial elements to take into account in the experimental design (McKay et al., 2023). Third, a multitude of animal studies have demonstrated that BPA impacts multiple reproductive outcomes, even in spite of the limitations inherent in clinical trials (Cloughesy et al., 2019). Firstly, in contrast to humans, rodents permit tissue- and cell-specific genetic analysis in addition to histological evaluation of a variety of tissues. Secondly, rodent models offer exceptional chances for sex-specific assessments (Edmonds et al., 2020). Important aspects that need to be taken into account in the experimental design include the tissue under examination, the structural, functional, and epigenetic changes between gender and species, the duration of exposure, and the dosage of the chemical (Jeremias et al., 2020). Third, numerous animal studies have demonstrated that BPA has an impact on multiple reproductive outcomes, even in spite of the limitations of clinical trials. Important aspects that need to be taken into account in the experimental design include the tissue under examination, the structural, functional, and epigenetic changes between gender and species, the duration of exposure, and the dosage of the chemical (Jeremias et al., 2020).

Third, numerous animal studies have demonstrated that BPA has an impact on multiple reproductive outcomes, even in spite of the limitations of clinical trials (Peretz et al., 2014). Oxidative stress and similar indicators are linked to BPA toxicity in a number of animal models, as well as in people (Meli et al., 2020). Growing research indicates that BPA organ damage is largely caused by the generation of Reactive Oxygen Species (ROS) and a reduced ability to defend against antioxidants, which changes the oxidative equilibrium in the mitochondria and throughout the cell (Amjad et al., 2020). In order to determine that oxidative stress is the leitmotif underpinning BPA's mechanism of action with pleiotropic effects on reproduction, this review provides fresh insights and an up-to-date literature scan on these topics (Burton & Jauniaux, 2011). Additionally, the study explores the impact of antioxidant compounds on BPA-induced toxicity and evaluates their potential effectiveness as a therapeutic approach to mitigate harm to the male and female reproductive organs and their functions (Sánchez et al., 2021).




Fig. 1: Mechanisms Connecting Reproductive Toxicity and BPA-Induced Oxidative Stress.




Fig 2: Protective Effects of Antioxidants on BPA Toxicity in Reproductive Systems.

Oxidative Stress as a BPA Toxicity Marker

Important for inflammatory responses, oxidative stress has been linked to immunological, cardio-metabolic, and aging disorders, neuro degeneration, and the onset and spread of cancer through a variety of unclear processes (Spahis et al., 2017). For cells to remain healthy, develop, grow, and survive, the redox balance must be regulated (Zhang et al., 2011). Superoxide anions, peroxides, and hydroxyl radicals (ROS) are produced by healthy cellular metabolism and are engaged in the well-regulated process

of redox balancing (Kumar & Pandey, 2015). Important for inflammatory responses, oxidative stress has been linked to immunological, cardio-metabolic, and aging disorders, neuro degeneration, and the onset and spread of cancer through a variety of unclear processes (Vallese, Cordone, Pecorelli, & Valacchi, 2023). For cells to remain healthy, develop, grow, and survive, the redox balance must be regulated (Ursini et al., 2016). Superoxide anions, peroxides, and hydroxyl radicals (ROS) are produced by healthy cellular metabolism and are engaged

in the well-regulated BPA is a metabolic and endocrine disrupting chemical that can impair oxidative homeostasis through direct or indirect mechanisms, such as increasing oxidative mediators and reducing antioxidant enzymes, determining mitochondrial dysfunction, changing cell signaling pathways, and inducing apoptosis (Ursini et al., 2016).

BPA induces oxidative stress by increasing lipid peroxidation and hydrogen peroxide in the kidney, brain, and testis of mice, and by decreasing antioxidant enzymes superoxide dismutase (SOD), catalase, glutathione reductase (GR), and glutathione peroxidase (GSH-Px) and decreasing antioxidant enzymes in the liver and epididymal sperm of rats related process of redox balancing (Kini et al., 2024). The ROS increase induced by BPA has been reported in several cell types with concentrations ranging from non-molar to micro molar (Yin et al., 2017). Differences in ROS generation, duration, and cytotoxicity appear mainly based on the concentration of BPA, on cellular backgrounds, exposure time, and sensor fluorescent reagents (Huang et al., 2022). BPA has been recently associated with inflammation markers in addition to oxidative stress in humans Yang and colleagues previously carried out a cross-sectional investigation involving men and women going through menopause or post menopause (Choi et al., 2017). They found that urinary BPA levels were positively correlated with urinary oxidative stress and blood inflammatory biomarkers (changes in white blood cell count and C-reactive protein) in women going through post menopause, suggesting that the effects of BPA vary depending on gender and hormonal status (Ferguson et al., 2016). According to these scientists, the effect of BPA toxicity on gender could be attributed to receptor occupancy and estrogen levels (Sifakis et al., 2017). The ER expression is known to be age and gender related. Due to the extremely low hormone levels in postmenopausal women, xeno estrogen BPA can bind to the ER more extensively, which can cause harmful cellular reactions like inflammation and oxidative stress (Matarrese et al., 2021). Oxidative stress and inflammation have a well-established association; they are closely related to cellular functions and may be the causes or effects of cell damage (Keane et al., 2015). The production of reactive oxygen species (ROS) and nitrogen species by immune cells such as neutrophils and macrophages has a role in the development of chronic inflammation (Fialkow et al., 2007). Widely regarded as potential biomarkers for oxidative stress, malondialdehyde (MDA) is a sign of lipid peroxidation, and 8-hydroxy-deoxyguanosine (8-OHdG) is a marker of DNA oxidation (Graille et al., 2020). Several investigations have shown their rise in BPA-exposed organisms (Ribeiro et al., 2017). The vicious loop between inflammation and oxidative stress is maintained by numerous different redox-sensitive signal transduction pathways, including c-Jun N-terminal kinase (JNK), p38 mitogen-activated protein (MAP) kinase, and transcription factor AP-1 (Sarg et al., 2024).

BPA-induced oxidative stress has been linked to the cytotoxicity of human B cells [30], the suppression of immunological response in vitro in murine macrophages, and the impairment of immune response in animal models

(Pirozzi et al., 2020). Additionally, exposure to BPA increases the expression of pro inflammatory cytokines in numerous tissues and organs (such as serum, the colon, the liver), as well as interleukin (IL)-1 β , IL-6, IL-8, and tumor necrosis factor- α (TNF- α) (Wang et al., 2019). Moreover, NOD-like receptor protein 3 (NLRP3) in acrosomes with inflammatory co-stimulus (i.e., nitric oxide) in adipocyte cells or the liver of BPA-exposed obese mice are activated in part by BPA-induced oxidative stress (Chen et al., 2023).

Mitochondrial impairment is one of the mechanisms of oxidative stress caused by BPA, behind lipid peroxidation and inflammatory reactions (Nagarajan et al., 2024). Low levels of BPA cause hepatic toxicity by affecting the liver's mitochondrial activity (Khan et al., 2016). Additionally, this pollution causes changes in the expression of genes related to mitochondrial activity and metabolism, ATP depletion, the release of cytochrome c, loss of mitochondrial mass and membrane potential, and mitochondrial metabolism (Sun et al., 2022). Sex differences in oxidative stress depend not only on the higher antioxidant defense in females but also on the increased generation of ROS in males (Tiberi et al., 2023). In fact, some studies have reported a gender-dependent NADPH oxidase activity, resulting in the excessive formation of reactive intermediates in male mice. Based on the intriguing literature concerning this issue, it will be relevant to carry out comparative studies on gender-related (Rosa & Flores, 2017).

Mechanisms Connecting Reproductive Toxicity and BPA-Induced Oxidative Stress

Males and Females have higher levels of reduced glutathione than males possess, and oxidative damage to mitochondrial DNA is lower in females than in males.

BPA Genetic Harm

The primary causes of genetic DNA damage are chromosomal abnormalities, which include aneuploidy, DNA strand breaks, DNA adducts, and gene mutation (Phillips & Arlt, 2009). Despite the fact that BPA is widely distributed and that people are constantly exposed to other contaminants when they are around, no information has been released regarding how BPA directly damages DNA by causing oxidative stress and the subsequent consequences this has on reproduction (Hengstler et al., 2011). Preclinical research conducted in vitro and in vivo has demonstrated that BPA induces mutations, aneuploidy, and DNA adduct formation (Tiwari et al., 2017). These changes could be a factor in the development of birth abnormalities, miscarriages, and infertility (Agenor & Bhattacharya, 2015). Additionally, BPA inhibits the base excision DNA repair mechanism that is brought on by laser irradiation (Gassman et al., 2015). The primary genetic abnormalities are chromosomal aberrations, which include aneuploidy, DNA strand breaks, DNA adducts, and gene mutation (Rejhová et al., 2018). Through the generation of MDA, phenoxyl radicals, and ROS, BPA-induced DNA damage has been linked to oxidative stress. This can result in direct DNA changes that are amplified by the disruption of antioxidant enzyme pathways (Dehghani et al., 2023).

Table 1: Classification of Antioxidant Approaches Against BPA Toxicity on Male and Female Reproductive Systems

Classification	Type	Dose (Suggested Values)	Notes
Vitamins and Co-Factors	Glutathione + Vitamin E	10 μ M + 5 mM	This combination counters BPA-induced oxidative damage and improves reproductive health in animal studies.
	Vitamin C + Vitamin E	100 mg/kg + 50 mg/kg	Co-administration enhances protection against oxidative damage in the ovaries and improves fertility.
	Vitamin E	6 mg/100 g bw	Supports testicular and sperm health, reducing BPA-induced apoptosis.
	1,25-Dihydroxyvitamin D ₃ (1,25D ₃)	0.2 μ M	Mitigates mitochondrial DNA damage and supports reproductive tissue health.
	Coenzyme Q10	200 mg/L	Enhances ROS reduction and prevents mitochondrial dysfunction in BPA-exposed models.
Natural Antioxidants	α -Lipoic Acid (LA)	30 mg/kg	Reduces oxidative stress and supports antioxidant enzyme activity.
	Retinoic Acid	10 mg/kg	Improves oxidative balance and reduces BPA's estrogenic effects.
	Gallic Acid	40 mg/kg	Counters lipid peroxidation and oxidative stress due to chronic BPA exposure.
	Cordyceps Militaris Extract	300 mg/kg	Enhances antioxidant defenses and reduces oxidative stress markers in reproductive tissues.
Hormones	Cuscuta Chinensis Flavonoids (CCF)	60 mg/kg	Blocks transcription of apoptotic proteins, improving sperm quality and reducing ROS levels.
	Melatonin	15 mg/kg	Reduces ROS and lipid peroxidation, while protecting mitochondrial integrity in BPA-exposed cells.
	Selenium	0.7 mg/kg	Reduces ROS and improves antioxidant defense mechanisms in reproductive tissues.
Trace Elements	Selenomethionine	4 mg/kg	Prevents testicular toxicity and enhances antioxidant enzyme activity.
	Betaine	5 g/kg	Supports maternal and neonatal health, reducing oxidative stress during BPA exposure in gestation.
Methyl Donors	Vitamin B ₁₂ + Folic Acid	20 μ g/kg + 20 mg/kg	Improves fetal development and reduces placental oxidative stress during gestation.

When *Drosophila melanogaster* was exposed to BPA through diet, found increased generation of reactive oxygen species (ROS) and lipid peroxidation, indicating impaired antioxidant function and mutagenesis consequences (Anet et al., 2019). Furthermore, the BPA treatment increased ROS generation in pig embryos, which resulted in cytochrome c release, damage to the mitochondria and DNA, and apoptosis via the p53-p21 signaling pathways (Mathur et al., 2012). After BPA was exposed to rat ovaries *ex vivo*, Ganesan et al. observed that BPA alkylating metabolite also caused damage to DNA (Ganesan & Keating, 2016). Urinary BPA levels were found to be correlated with MDA and/or 8-OHdG in several human studies (Steffensen, 2020). BPA blood levels were found to positively correlate. Prostate epithelial cells also showed signs of BPA genotoxicity: following a 24-hour exposure to escalating doses, BPA suppressed p53 expression (Evans, 2022). The prostate cells treated with high or low concentrations of BPA had DNA adducts (De Flora et al., 2011).

The same authors also found that when male rats were exposed to BPA (200 mg/kg for 10 days), there was a noticeable increase in MDA and nuclear spreading factor in spermatozoa, which may indicate that these germ cells are fragile (Sadek et al., 2024). In spermatogonia (GC-1 cell line), micromolar BPA doses have been shown to cause ROS generation, DNA damage, and cytotoxicity. Remarkably, at non-cytotoxic micromolar and nanomolar doses, the induction of DNA strand breaks and its association with ROS production were also noted (Meli et al., 2020). Other cell lines, however, did not exhibit the induction of strand breakage by non-cytotoxic doses of BPA. The differing views about the genotoxic effects of

BPA on DNA may be due to variations in BPA concentrations used, which are frequently higher than those found in ambient matrices. BPA causes genotoxic effects in somatic and germ cells when tested *in vitro*, but it cannot be regarded as a "true" genotoxic agent *in vivo* because it does not cause cancer in animals (Bariar, Vestal, & Richardson, 2013). It is unclear if BPA can have an indirect genotoxic effect in the context of other environmental contaminants or pathological circumstances, as it cannot induce genetic damage when present in environmental amounts (Langie et al., 2015).

Effects of BPA Epigenetics

With distinct patterns that change based on the kind of cell, epigenetic mechanisms control gene expression and cellular phenotypes without changing the nucleotide sequence of the genes (Chen, 2007). Numerous investigations have demonstrated the detrimental effects of epigenetic modifications and DNA damage on early embryonic development (Ornoy, Reece, Pavlinkova, Kappen, & Miller, 2015). The DNA methylation and histone modification patterns undergo significant programming and reprogramming throughout this time, and the epigenetic mechanisms can modify the male and female genomes in a tissue-specific manner through activation/deactivation processes (Navarro et al., 2020). Methylation limits the binding of transcription factors to DNA, which is crucial for mammalian development, or it induces gene silence by proteins involved in gene expression (Ehrlich & Lacey, 2013). According to published research, the epigenetic effects of BPA and other environmental contaminants on reproduction vary based on the type of exposure (dose, duration, and route),

the gender, and the stage of life (Mileva et al., 2014). BPA was given to female agouti yellow mice at a dose of 50 mg/kg by food, causes a DNA hypomethylation that results in phenotypic changes in the coat color of offspring (from the mating to the weaning stage) (Rosenfeld et al., 2013).

On the other hand, Doshi et al. found that BPA administration caused hypermethylation of estrogen receptor promoter regions in the testicles of newborn rats during the first five days of life, which interfered with spermatogenesis and reduced fertility (Cariati et al., 2020). BPA exposure, beginning at low concentrations, may influence both DNA hypermethylation and hypomethylation in CpG islands during fetal development and in a dose-dependent way (Onuzulu et al., 2019). Contradictory findings from other studies indicate that BPA-induced epigenetic modifications have the opposite effect on DNA methylation; the underlying molecular and cellular mechanisms are yet unknown. Indeed, chromosomal disintegration with both DNA hypermethylation and hypomethylation has been linked to increased ROS generation and oxidative stress (Chianese et al., 2018). BPA has been shown to cause epigenetic reproductive harm, and oxidative stress and sperm epimutations may be epigenetically passed down to progeny through DNA methylation (Lombó et al., 2019). Oral exposure to BPA during gestation and/or lactation in pregnant albino rats resulted in increased MDA levels and DNA hypermethylation at CpG islands in the promoter genes of ER α and DNA methyltransferases in the testis of the male offspring (El Henafy, Ibrahim, Abd El Aziz, & Gouda, 2020). There was also a description of the decline in superoxide dismutase (SOD), GST, and GSH-Px levels (Chattopadhyay et al., 2007).

This data aligns with our unpublished findings about BPA-induced liver metabolic impairment in obese mice. The global DNA hypermethylation (% of methyl cytosine/total) brought on by high-fat diet is made worse in our experimental setting by a 21-day oral BPA exposure (50 μ g/kg daily in corn oil) (Martos et al., 2017). According to an *in vitro* investigation, acute BPA caused aberrant cytoskeletons, lower maturation rates, elevated ROS generation, DNA methylation, and apoptosis/autophagy rates in pig oocytes (Sharma, Sharma, Gupta, & Srivastava, 2019). The authors came to the conclusion that BPA might cause additional disruptions to oocyte meiosis by altering the state of DNA methylation. After exposing pig embryonic-derived parthenoides to BPA, Guo et al. found that increased ROS production was linked to mitochondrial and DNA damage, which in turn caused autophagy and altered DNA methylation (decreasing 5-methyl-cytosine and DNA methyl transferase RNA) and may have an impact on the health of the offspring (Anderson et al., 2001). On the other hand, the impact of a subacute BPA exposure on an adult male minnow's testes, the rise in the generation of hydrogen peroxide antioxidants and the hyper methylation of DNA worldwide (Santiago et al., 2021).

Metabolic and Endocrine Disruption

By changing hormone production and metabolism and binding to hormone receptors, endocrine disruptors can

alter hormonal balance (Nadal, Quesada, Tuduri, Nogueiras, & Alonso-Magdalena, 2017). These changes can lead to a wide range of developmental and reproductive problems (Skakkebaek et al., 2016). Numerous studies conducted recently, including *in vitro*, *in vivo*, and epidemiological ones, have connected EDC exposure to a higher risk of developing a number of diseases, including obesity, metabolic syndrome, type 2 diabetes, and cardiovascular diseases (Lind, Ingvaldsen, & Furevik, 2018). In fact, some disruptors really cause metabolic illnesses directly, while others work by raising the disease's threshold or sensitivity (Casals-Casas & Desvergne, 2011). There is ample evidence that it is difficult to objectively evaluate the dangers associated with EDC exposure due to the variety of chemical structures and methods of action, as well as factors such as developmental stage, timing, and dose exposure (Fuhrman, Tal, & Arnon, 2015). BPA, mimics the structure and actions of 17- β -estradiol; additionally, it binds to GPR30 and other conventional and non-classical membrane estrogen receptors (Molina-López et al., 2023). These activities cause various alterations in the ovary, mammary gland, and uterus organs that target estrogen in females exposed to BPA during pregnancy (Paulose et al., 2015).

Furthermore, BPA suppresses receptor degradation and ubiquitination by interfering with target gene transcription that is mediated by ER β (Amir et al., 2021). Moreover, BPA functions as an antagonist by binding to the androgen receptor. It is well known that the androgen receptor plays a key regulatory role in androgen cell signaling and is crucial for the development and function of male reproduction (Perera et al., 2017). Together with thyroid hormone activity, the hypothalamic-pituitary-testicular axis regulates both human and animal spermatogenesis (La Vignera & Vita, 2018). The correlation between BPA exposure and lower levels of testosterone in Chinese men as well as higher serum levels of luteinizing hormone (LH) and follicle-stimulating hormone (FSH) (X. Liu et al., 2015). As is well known, FSH and LH promote testosterone synthesis and boost prostate growth by acting on Sertoli and Leydig cells, respectively (Ramaswamy & Weinbauer, 2014). According to a recent review, BPA has a negative impact on male reproductive hormones and has comparable effects on thyroid hormones, which are known to affect sperm production and male fertility (Cariati et al., 2020). Thyroid hormone levels were shown to be lowered in newborn girls exposed to BPA later in gestation, and thyroid illness may be associated with BPA-induced ROS generation (Mohammed et al., 2020).

The connection between BPA and metabolic illnesses (such as obesity and diabetes) can have an impact on both male and female fertility and reproduction by causing fat buildup and impairing the metabolism of fats and carbohydrates, which is linked to inflammation (Heindel et al., 2017). BPA is a lipophilic compound that builds up in fat, increasing the quantity and size of adipocytes and causing weight gain. BPA, as described before, modifies the release of adipokines at ambient levels by decreasing anti-inflammatory adiponectin and increasing pro-inflammatory leptin and other inflammatory cytokines

(e.g., IL-6, TNF- α) (Ahmed, 2020). It is well known that the relationship between energy balance and reproduction depends on adipokines like resistin and leptin as well as gastrointestinal peptides (Comminos et al., 2014). Similar to insulin resistance, high levels of leptin are associated with fat mass and leptin resistance, which are common in obesity and related diseases such as diabetes (Zimmet et al., 1999). It's interesting to note that people who have greater blood levels of BPA are more than twice more likely to acquire diabetes than people who have lower levels of BPA, and that risk increases with age (Hwang et al., 2018).

Signaling Cell:(Zimmet, Boyko, Collier, & de Courten, 1999)

Depending on the type and function of the cell, BPA activates one or more distinct and numerous cell signaling pathways (Murata & Kang, 2018). In non-genomic signaling, BPA binds to GPR30 and mER, activating these receptors and triggering rapid estrogenic signaling that activates kinase systems, including phosphoinositide 3-kinases (PI3K), protein kinase-A (PKA), and MAPK (Ahmad et al., 2024). It also modifies intracellular calcium levels in cancer cells, cAMP, and protein kinase-C (PKC) (Kamp & Hell, 2000). According to a recent summary, spermatozoa exposed to BPA in vitro activate these receptors, which are linked to elevated ROS (Cariati et al., 2020). While ROS are necessary for optimal sperm activity at low levels, large amounts of ROS cause damage to cellular lipids, proteins, and DNA as well as adverse effects on sperm motility, mobility, and acrosome reaction (Dutta et al., 2019). These changes quickly lead to increased ROS availability in the cytoplasm and activation of MAPK, PI3K, and PKA (Krylatov et al., 2018). Increased production of reactive oxygen species (ROS) leads to phosphorylation of sperm proteins in their tyrosine residues, which in turn causes functional modifications of core regulatory genes and proteins through transcriptional and translational pathways, hence predisposing cells to aberrant outcomes (Shah & Brownlee, 2016).

Hence, BPA may impair spermatozoa's complicated oxidative stress response, which may have an impact on their capacity to fertilize (Ryu et al., 2023). It has also been shown that BPA causes increased ROS and lipid peroxidation in various cell types (Huc et al., 2012). It had been reviewed that BPA affects many cell-signaling mechanisms and pathways extracellular signal-regulate kinases (ERK), JNK involve dinflammation, immune response, cancer, brain, and reproductive function; the increase of calcium and ROS and the activation of ERK and JNK are implicated in BPA-induced apoptotic cell death Several studies have determined that EDC-induced oxidative stress can impair male fertility by disrupting the cell junctions among cell testis (Xiao et al., 2020). The pathways involved in BPA male reproductive toxicity, analyzing the pathophysiological role of MAPK and inflammatory cytokines and that of the PI3K/c-Src/ focal adhesion kinase signaling pathway (Wong, Boon-Itt, & Wong, 2011). The resulting alteration of intercellular junctions, via polarity proteins, leads to reproductive dysfunction such as reduced sperm count and reduced semen quality (Costa & Mercadal, 1995).

Oxidative Stress-Induced BPA Toxicity on Male and Female Reproduction

Many conflicting studies on the impact of BPA on male fertility were gathered between 2000 and 2010, although numerous additional recent findings have consistently shown that BPA has negative effects on both male and female reproductive processes (Mendiola et al., 2010). Transgenerational investigations have shown that BPA specifically affects spermatogenesis and sperm quality in males; it also affects ovary, embryo development, and egg cell quality in females (Rizzo et al., 2023). In addition to its direct effects on reproductive organs, BPA also affects the ability of the offspring to reproduce because of its ability to penetrate the blood brain barrier (Yahav et al., 2019). Reproductive function changes have a permanent impact on adult fertility, particularly if they happen while the reproductive organ is still developing in a fetus especially oxidative stress (Al-Gubory et al., 2010). Numerous preliminary results have demonstrated the influence of perinatal BPA exposure on male spermatogenesis through various modes of action (Lombó et al., 2019).

That oral BPA exposure in gestational day (GD) dams enhanced generation of reactive oxygen species (ROS) and activated the apoptotic pathway in the testis of male offspring, as shown by histological alterations (Quan et al., 2017). In addition to changing sperm quality and motility in adult male mice, BPA treatment in CD-1 dams (GD7–14) also affected oxidative balance. Additionally, impaired spermatogenesis was seen in progenies in a mouse model of chronic BPA exposure of dams throughout the entire gestational period and continuing the oral delivery to male pups until sexual maturation (Delbes et al., 2022). This impairment was linked to a decline in antioxidant defense as well as a decrease in the expression of sirtuin 1, a crucial ROS generation sensor (Singh et al., 2018). Data from very recently confirmed that male pups exposed to BPA at a young age developed oxidative stress in their testicular tissue (Sedha, Kumar, & Shukla, 2015). It was shown in this comparative study between prenatal, perinatal, and postnatal BPA exposure that BPA-related epigenetic modifications happen irrespective of temporal exposure (Kunysz et al., 2021).

However, the level of BPA's harmful impact on oxidative balance varied with period of exposure, peaking during perinatal exposure (from pregnancy to nursing) (Varshavsky et al., 2020). Adult rats exposed to BPA experienced a dose-dependent increase in reactive oxygen species (ROS) (Babu et al., 2013). Oxidative stress was linked to metabolic impairment (hyperglycemia and hyperinsulinemia) and reproductive damage, as evidenced by the testicular reduction of insulin receptor substrate-2 and glucose transporter-8, two important proteins involved in the metabolism of testosterone and spermatogenesis (Dias, Parente, Vasconcelos, & Dias, 2014). BPA caused a significant increase in lipid peroxidation in the epididymis, testis, and immune cells (i.e., lymphocytes and bone marrow) of adult rats (Srivastava et al., 2015). This increase is linked to sperm quality impairment and the induction of several antioxidant enzymes, including SOD, catalase, and non-enzymatic reduced glutathione (Micheli et al., 2016). In the testicles of offspring from dams exposed to BPA,

Khalaf et al. also recently discovered elevated levels of H₂O₂, lipid peroxidation, and depletion of antioxidant defense mechanisms (Marcu et al., 2023). BPA changed the shape of the seminiferous tubule epithelium, reduced gonadotropin discharges, increased ROS generation, and decreased the antioxidant activity of catalase and SOD enzymes during the early stages of puberty (Zahran et al., 2021). The primary cause of sperm failure and male infertility in the testis is increased oxidative stress, which is brought on by persistent alterations to the redox balance (Barati et al., 2020). These alterations result in ROS overproduction and the weakening of antioxidant defense (de Haan et al., 2003). It is crucial to emphasize that BPA's reproductive toxicity to oxidative stress was explicitly addressed in the clinical investigations (Bameri et al., 2024).

Yet, given the significance of oxidative balance in the physiology of sperm, oxidative stress may be the primary harmful mechanism responsible for BPA's toxicity to the male reproductive system (Ryu et al., 2023). The main factor contributing to male infertility and sperm failure in the testis is elevated oxidative stress, which is caused by ongoing changes in the redox balance (Bisht et al., 2017). The overproduction of ROS and the weakened antioxidant defense are the outcomes of these changes (de Haan et al., 2003). It is imperative to underline that the clinical studies specifically addressed BPA's reproductive toxicity to oxidative stress (Ahmad et al., 2024). Yet, given the role of oxidative balance in the physiology of sperm, oxidative stress may be the major damaging mechanism responsible for BPA's toxicity to the male reproductive system (De Toni et al., 2020).

Protective Effects of Antioxidants on BPA Toxicity in Reproductive System

Differently sourced chemicals known as antioxidants stop the oxidation or overproduction of reactive oxygen species (ROS) in cells and tissues, hence averting oxidative stress (Valko, Rhodes, Moncol, Izakovic, & Mazur, 2006). Many studies conducted in the past few decades have shown how antioxidants including gallic acid, lipoic acid, N-acetylcysteine, ginger extract, and vitamins C and E can protect or lessen the toxicity caused by BPA in a variety of organs (Li et al., 2022). The preservation of spermatozoa's quality, vitality, motility, and morphology is one of the benefits linked to antioxidant agent supplementation (Cilio et al., 2022). It's interesting to note that antioxidants maintain sperm motility and fertilization potential while reducing oxidative stress in spermatozoa during cryopreservation (Bansal & Bilaspuri, 2011). Here, we look at several antioxidant classes that, when taken singly or in combination, have been shown to be effective in reducing the harmful effects of BPA on the reproductive systems of both men and women (Santiago et al., 2021).

Vitamins and Co-Factors

It has been shown that vitamins C, E, and glutathione affect sperm function (Yousef, Abdallah, & Kamel, 2003). Specifically, these three drugs inhibited the excessive release of reactive oxygen species (ROS) and increased intracellular ATP synthesis in spermatozoa exposed to BPA, hence preventing motility loss and aberrant acrosome reactivity (Amaral et al., 2016). The primary

cause of the reduction in spermatozoa motility and fertilizing competence is increased ROS and their metabolites, which damage cells irreversibly by interfering with their enzyme systems, DNA, lipids, and proteins. Vitamin E (2 mM) and glutathione (5 mM) mitigated the effects of BPA (100 µM) on impaired fertilization and early embryo development (Dattilo et al., 2016). One of the main endogenous antioxidants, glutathione's recycling is influenced by vitamin C and E levels and vice versa (Bouayed & Bohn, 2010). In fact, ascorbic acid, or vitamin C, is an electron in addition to its scavenging action (Santos et al., 2022). In the ovaries of rats exposed to a high dose of BPA, the combination of vitamins C (50 mg/kg) and E (50 mg/kg), but not these vitamins alone, was also able to exhibit a protective effect in reducing apoptotic cell death (Bilgi et al., 2019). It is well known that male germ cells contain large levels of vitamin E, and that vitamin E is essential for controlling the lipid peroxidation that takes place in testicular microsomes and mitochondria (Ourique et al., 2016). There is proof that male fertility was enhanced by vitamin E (4 mg/100 g bw), which shielded testicular cells and epididymal sperm from the apoptosis caused by BPA exposure in Wistar rats (Justin et al., 2024). Recent research has shown that 1, 25-dihydroxyvitamin D₃ (1,25D₃; 0.1 µM) can modify the harmful effects of BPA (10 µM) on oxidative stress, specifically on the dynamics and function of mitochondria in ovarian granulosa cells (Jain et al., 2024). Furthermore, 1,25D₃ shielded the female ovary against BPA-induced mitochondrial DNA loss (P. Liu et al., 2023). It's interesting to note that women who have been exposed to BPA had lower serum vitamin D levels (Brennan et al., 2024). Coenzyme Q10 (CoQ10), among other vitamins, has the ability to restore fertility and mitigate the reproductive toxicity linked to BPA exposure (500µM) in the *Caenorhabditis elegans* germ line (Carneiro et al., 2023).

By reducing oxidative stress through the scavenging of ROS and free radicals, decreasing meiotic DNA double-strand breaks, and restoring mitochondrial activity, CoQ10 partially counteracted BPA-induced DNA damage (Carneiro & Colaiácovo, 2023). In the context of BPA-treated worms, CoQ10 supplementation resulted in a decrease in the number of aneuploid embryos and chromosome abnormalities in oocytes at diakinesis (Bradford, Schlaepfer et al., 2020). All prokaryotic and eukaryotic cell types contain the biological thiol LA, which plays a crucial function in mitochondrial damage and lowers oxidative stress while also raising the amounts of other antioxidants (Herrero et al., 2008). As a biofactor of energy metabolism, LA responds powerfully to reduce oxidative stress and free radicals (André et al., 2024). It also prevents the decline in glutathione levels, restricts the production of lipid peroxides (LPOs), and restores the balance of antioxidant enzymes in the body (Marí et al., 2009). In male rats, prolonged BPA (10 mg/kg) administration resulted in testicular and mitochondrial oxidative stress (Anjum et al., 2011). To be more precise, LA boosted both enzymatic and non-enzymatic antioxidants of mitochondria, such as glutathione reductase and succinate dehydrogenase, malate dehydrogenase, isocitrate dehydrogenase, monoamine oxidase, and NADH dehydrogenase, and corrected BPA-altered

activity of these essential mitochondrial enzymes (Bayliak et al., 2023).

Ultimately, oral LA treatment (100 mg/kg) for 30 days during BPA exposure (25 mg/kg, per os) in female Wistar rats resulted in a reduction of ovarian oxidative damage, preventing nitrosative stress and lipid peroxidation, and even more in combination with vitamin E (20 mg/kg, per os) (Carneiro & Colaiácovo, 2023). Retaining retinoic acid, a metabolite of vitamin A, may also protect against oxidative stress damage was found to increase the expression of genes linked to antioxidants in mature buffalo oocytes cultured in vitro (Rakha et al., 2022). Furthermore, in both healthy and varicocele sperm, all-trans retinoic acid increased glutathione transferase and SOD activities while lowering reactive oxygen species and MDA levels, indicating that retinol increases the activity of antioxidant enzymes (Malivindi et al., 2018). According to Koda et al., adult ovariectomized rats supplemented with 5 mg/kg of all-trans retinoic acid per os greatly inhibited the rise in uterine weight caused by BPA (100 mg/kg), hence blocking the estrogenic activity of BPA (F. Amjad et al., 2021). It's interesting to note that vitamin A-containing pomegranate juice reduces sperm abnormalities and increases caudal sperm count in rats by preventing BPA-induced structural alterations in the epididymis whereas hepatic retinoids are necessary for BPA biotransformation and subsequent (Funk & March, 1897).

Natural Extracts and Products

Among several preventive or therapeutic methods, it is advisable to contemplate the utilization of natural substances like phytochemicals and herbs (Ahmad et al., 2024). Mostly found in plants, fruits (such as pineapple, banana, grapes, gallnuts, apples, and lemons), and vegetables, gallic acid is an apolyhydroxyphenolic molecule that can either be free or partially hydrolyzed tannin (Kang et al., 2018). This chemical has been shown to have antioxidant qualities, which are associated with scavenging ROS and reserving metal ions (Balogh et al., 2003). Gallic acid (20 mg/kg in maize oil) has recently been found to have a role in testicular oxidative stress in adult rats exposed to chronic BPA (10 mg/kg) (Ola-Davies & Olukole, 2018). These authors demonstrated that Gallic acid restored the BPA-altered parameters of antioxidant enzymes, reduced testicular lipid peroxidation (MDA, MPO), and normalized the gonadosomatic index (which is used to assess sexual maturity in connection to testicular development) (Gezer et al., 2024). Cordyceps militaris is a fungus that is used in traditional Chinese medicine to treat impotence, seminal emission, and infertility (Bach et al., 2022). It exhibits a variety of biological activities, including an antioxidant impact because of its polysaccharide and cordycepin content (Jędrejko et al., 2021).

The preventive effect of Cordyceps militaris extract (200–800 mg/kg) against the reproductive harm caused by BPA poisoning (200 mg/kg) in rats has been studied (Salim et al., 2023). The natural product's dose-related effect led to an increase in sperm count and motility, which were both affected by BPA (Sharma et al., 2019). It also reduced oxidative stress and the associated alterations in adipocyte function. GSH levels and antioxidant enzyme activity

were found to be enhanced by these effects (Peris et al., 2019). The primary constituents of Chinese herbal medicine *C. chinensis*, called *Cuscutachinensis* flavonoids (CCF) have anti-abortion and antioxidant properties (Hajimehdipoor et al., 2012). It has been found recently that these flavonoids (40 mg/kg) can lessen the testicular cell apoptosis caused by BPA (5 mg/kg) (Zhao et al., 2023). This effect in male mouse offspring (F1) following contextual maternal exposure to BPA and flavonoids is directly linked to oxidative damage. Specifically, in the testes of F1 mice collected at postnatal days 21 and 56, CCF inhibited the transcription and translation of apoptotic proteins (caspase 7 and 9), thereby minimizing structural damage and improving fertility (Udeochu et al., 2023). Recently, mice given BPA (1 mg/kg) orally showed protective effects from hydroethanolic *Murrayakoenigii* extract of leaves (200 mg/kg) (Kaur & Gupta, 2020).

Melatonin

Melatonin is a neuro-hormone that is mostly secreted from the pineal gland. It is generated from tryptophan and operates on the hypothalamic-pituitary-gonadal axis (Mocchegiani et al., 1994). Melatonin is engaged in various homeostatic processes, such as controlling reproduction, maintaining circadian rhythms, and maintaining redox balance (Megha et al., 2024). Numerous experimental and clinical investigations emphasizing the safety of melatonin have led to a significant increase in interest in this substance as an antioxidant (Ramos et al., 2021). In fact, melatonin functions as an inducer of antioxidant enzymes and a scavenger of various nitrogen and ROS species, such as hydroxyl radicals, peroxy nitrite anions, and H₂O₂ transcription of an enzyme (Tan et al., 2007). Because of its molecular makeup, melatonin can pass through cellular membranes and enter the cytosol, mitochondria, and nucleus, protecting DNA and enhancing signaling (Kopustinskiene & Bernatoniene, 2021). A prior study demonstrated the protective effect of melatonin (10 mg/kg, i.p.) on adult mice's interstitial mitochondrial damage caused by brief exposure to BPA (10 mg/kg) (Priego et al., 2021). Additionally, following a prolonged exposure, melatonin (10 mg/kg, i.p.) was shown to be able to reverse BPA (50 mg/kg)-induced oxidative damage and apoptosis in rats' testes and the change of epididymal sperm (Sahu & Verma, 2023).

Melatonin combined the anti-apoptotic effect on rat germ cells with its capacity to regulate cellular redox balance (Abo El Gheit et al., 2022). In fact, melatonin decreased apoptosis by increasing the redox-sensitive and anti-apoptotic protein Bcl-2, and it also positively regulated glutathione, SOD, and catalase, as well as MDA levels and H₂O₂ output in the tissues and sperm (Lu et al., 2018). Due to its manipulation of blood testosterone levels, melatonin's protective action on spermatocyte meiosis and proper sperm quality and production is also strongly linked to its control of cellular redox balance (Loren et al., 2017). Since the effectiveness of melatonin supplementation was shown to affect DNA alterations, it has been established that the hormone (10 mg/kg) enhanced the genotoxicity potential of BPA (200 mg/kg), lowering the amount of damaged DNA in adult rats' male germ cells by suppressing oxidative

stress (Mączka, Grabarczyk, & Wińska, 2022). Specifically, spermatocyte immunostaining revealed that the melatonin preventive action reduced DNA movement within germ cells and the formation of γ H2AX, a hallmark of DNA double-strand breaks (Geesink & Meijer, 2017). Alongside this effect, melatonin-induced decreases in the levels of reactive chemicals thiobarbituric acid and elevations in SOD were observed (Essawy et al., 2025). The protective impact of melatonin on adrenal and prostate function in adult male rats exposed to oral BPA for two weeks (Olukol et al., 2019). Later, after mother exposure to various levels of BPA, further investigated the pertinent effect of melatonin (1 mg/kg) in shielding F1 adult rats against male reproductive damage (Tain & Hsu, 2024). In addition to improving testicular SOD, GSH, and GPx activity, melatonin's antioxidant-protective effect on interstitial necrosis in testis (caused by BPA administration in utero) was linked to limiting germinal cell degeneration and recovering BPA-altered tubular and luminal diameter. The beneficial effects of melatonin (20 mg/kg) on the changes in epididymal structure and sperm quality brought on by long-term exposure to BPA (25 mg/kg) (Heidarizadi, Rashidi, Jalili, & Gholami, 2022). There have also been reports of BPA having negative effects on the female reproductive system (Matuszczak et al., 2019). The mechanisms behind the deleterious effects of BPA (100 μ g/kg bw per day for 7 days) on mice's oocyte quality (meiotic maturation and fertilization capacity) (Yahav et al., 2019). By lowering ROS levels and preventing oocyte apoptosis, the scientists found that oral melatonin (30 mg/kg bw per day for seven days) increased the in vitro fertilization rate and restored BPA-induced changes to fertilization proteins and processes (M. Zhang et al., 2017). Additionally, melatonin guards against BPA-induced uterine deterioration (Bameri et al., 2024).

Selenium

The selenium plays a crucial function in reducing spermatogenic oxidative damage and apoptosis as well as preserving male fertility after BPA administration (Sahu & Verma, 2023). Consequently, selenium's capacity to regulate the crucial balance between germ cell death and proliferation during apoptosis in spermatogenesis has been previously proven (Yan et al., 2024). Selenium supplementation in the diet (0.5 mg/kg diet) decreased ROS and lipid peroxidation in mouse testes and ameliorated histopathological abnormalities, conserving basement membrane and lowering germ cell vacuolization in mice exposed to BPA (150 mg/kg) (Mondal & Bandyopadhyay, 2024). The interaction between selenium and selenoproteins, such as the ROS-interacting enzyme GSH-Px, is most likely the cause of this impact (Salvador et al., 2020). In male rats exposed to 150 mg/kg of BPA, a recent study verified the protective effects of selenium (3 mg/kg) and nanoselenium (2 mg/kg) against reproductive damage. The selenium nanoparticle known as nanoselenium has lower toxicity, more surface activity, and better bioavailability than regular selenium (Zhou et al., 2019). Both elements have been shown to have a protective effect against BPA-induced testicular toxicity in rats; they also improved oxidative stress biomarkers, DNA fragmentation damage, inflammation, and the

expression of particular spermatogenesis-related genes (Khalaf et al., 2019).

Methyl Donors

Rats' BPA-induced unfavorable metabolic impact and oxidative stress were discovered to be improved, among other therapeutic techniques, by treating N-acetylcysteine (Elswefy et al., 2020). In the sulfation cycle, cysteine participates by serving as a donor of sulfur and a methyl donor during the conversion of homocysteine (Hcy) to methionine (Elswefy et al., 2020). Auto-oxidation of hcy produces reactive oxygen species (ROS) and may be involved in the beginning of reproductive problems. Hcy induces intrauterine abortion, premature birth, and low birth weight offspring when it is present in high amounts in the serum of pregnant women (Cetin et al., 2010). Supplementing the diet of their offspring with methyl donors at doses capable of lowering hemoglobin levels in plasma alleviated intestinal digestion and absorption disorders brought on by BPA, which may be linked to DNA methylation (Mączka et al., 2022). Sows have shown that supplementing with vitamin B9, B6, B12, and betaine lowers plasma Hcy levels and improves reproductive function (Rizzo et al., 2023).

Consistent with these results, have recently shown that methyl donor supplementation in the mother's diet (3 g/kg betaine, 400 mg/kg choline, 150 μ g/kg vitamin B12, and 15 mg/kg folic acid) during gestation effectively counteracted oxidative stress caused by BPA in the placenta and helped to maintain a healthy redox environment in the pig offspring as well as the fetus (Mou et al., 2018). The detrimental effects of BPA on placental integrity were actually mitigated by a dietary methyl donor, which also increased the antioxidant activity of enzymes (SOD, catalase, and GSH-Px) acting on placental trophoblastic cells and umbilical cord blood (Feng et al., 2023). These findings then had an impact on the maintenance of the redox balance in newborn piglets. It's interesting to note that how mother nutrition affected the properties of the muscles in the offspring following prenatal BPA exposure and co-treatment with methyl donors (Acharyya, 2007). BPA decreased methylation at the LDH promoter and increased the activity of the enzyme lactate dehydrogenase (LDH) as well as the amount of mRNA in the thoracic muscle at birth and during the finishing period (Ghimire et al., 2022). Through DNA methylation, the protective treatment with methyl donors has long-lasting effects on the quality of hog flesh by influencing the expression of glutathione and LDH (Nordberg et al., 2010).

Conclusions

Because BPA interferes with metabolism and endocrine function and has been linked to human diseases like cancer, diabetes, obesity, and reproductive issues, it has become a focus of significant investigation. This review offers a paradigm for comprehending how the pleiotropic reproductive deleterious effects following BPA exposure may be facilitated by the production of oxidative stress. Male reproductive organs are more susceptible to BPA exposure than female reproductive organs, according to multiple researches that have been reviewed. Research indicates that men may be more vulnerable for a number

of reasons: (i) gender differences in glutathione levels and their relevance for the detoxification process (males have lower glutathione availability); (ii) females have a greater capacity for sulfate-based detoxification; (iii) male reproductive organs have a greater inflammatory response; and (iv) females are less vulnerable to oxidative stress.

BPA exposure during infancy and development can cause oxidative stress and lipid peroxidation, which can affect hormonal and metabolic processes in the reproductive system and be conserved in both humans and animal species. Inflammation will often arise and intensify oxidative stress, and vice versa. Oxidative stress is frequently the initial anomaly in an organ. Growing and convincing evidence indicates that a wide range of BPA doses or concentrations (in vivo and in vitro or human exposure) promote ROS generation and alter redox balance, inducing mitochondrial dysfunction and the modulation of cell signaling pathways related to oxidative stress, despite differences among doses, duration, model systems, and measured outcomes. In addition to its endocrine and metabolic disrupting qualities, BPA can also generate oxidative stress, which can act in a dependent or independent manner.

This can have significant effects on reproduction throughout prenatal, perinatal, and postnatal exposure, as well as in adulthood. However, when BPA exposure is coupled with additional risk factors including a poor diet, impaired metabolism, and coexisting disorders, the effects can be quite severe. Various antioxidant strategies have been found to date to mitigate BPA-induced harm to the reproductive systems of men and women. By lowering certain oxidative stress markers, lipid peroxidation, or DNA damage, and reestablishing the antioxidant defense, these substances increase both male and female fertility. To determine the specific molecular processes underlying these antioxidant compounds' protective effect on oxidative stress and inflammation which are responsible for the systemic and organ-specific toxicity of this ubiquitous xenobiotic new experimental and clinical research are necessary. More research on antioxidants is required to determine the best dosage and course of action for addressing BPA poisoning as well as to better characterize their value in a number of gender-dependent reproductive dysfunctions.

DECLARATION

Funding: This study did not receive any national or international funds.

Acknowledgement: None.

Conflicts of Interest: All authors of the manuscript declare that they have no financial or personal interests.

Availability of Data and Material: All the data is available and can be obtained on reasonable request from the corresponding author

Ethics Statement: The research trial was conducted according to the guidelines regarding welfare and use of

animals for research purpose.

Authors Contribution: AN and NW: Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Software, Validation, Visualization, Writing-original draft, Writing - review and editing; NA, NA and SR: Data curation, Formal analysis, Investigation, Writing-original draft, Writing-review and editing.

Generative AI Statement: The authors declare that no Gen AI/DeepSeek was used in the writing/creation of this manuscript.

Publisher's Note: All claims stated in this article are exclusively those of the authors and do not necessarily represent those of their affiliated organizations or those of the publisher, the editors, and the reviewers. Any product that may be evaluated/assessed in this article or claimed by its manufacturer is not guaranteed or endorsed by the publisher/editors.

REFERENCES

- Abo El Gheit, RE, Soliman NA, Nagla SA, El-Sayed RM, Badawi GA, Emam MN, Radwan DA (2022). Melatonin epigenetic potential on testicular functions and fertility profile in varicocele rat model is mediated by silent information regulator 1. *British Journal of Pharmacology*, 179(13), 3363-3381.
- Abraham A & Chakraborty P (2020). A review on sources and health impacts of bisphenol A. *Reviews on environmental health*, 35(2), 201-210.
- Acharyya S (2007). Collisional emplacement history of the Naga-Andaman ophiolites and the position of the eastern Indian suture. *Journal of Asian Earth Sciences*, 29(2-3), 229-242.
- Agenor A & Bhattacharya S (2015). Infertility and miscarriage: common pathways in manifestation and management. *Women's health*, 11(4), 527-541.
- Ahmad I, Kaur M, Tyagi, D, Singh TB, Kaur G, Afzal SM & Jauhar M (2024). Exploring novel insights into the molecular mechanisms underlying Bisphenol A-induced toxicity: A persistent threat to human health. *Environmental toxicology and pharmacology*, 108, 104467.
- Al-Gubory KH, Fowler PA & Garrel C (2010). The roles of cellular reactive oxygen species, oxidative stress and antioxidants in pregnancy outcomes. *The international journal of biochemistry & cell biology*, 42(10), 1634-1650.
- Amaral S, S Tavares R, Baptista M, I Sousa M, Silva A, Escada-Rebelo S, Ramalho-Santos J (2016). Mitochondrial functionality and chemical compound action on sperm function. *Current medicinal chemistry*, 23(31), 3575-3606.
- Amir S, Shah STA, Mamoulakis C, Docea AO, Kalantzi OI, Zachariou A, Makriganakis A (2021). Endocrine disruptors acting on estrogen and androgen pathways cause reproductive disorders through multiple mechanisms: a review. *International Journal of Environmental Research and Public Health*, 18(4), 1464.

- Amjad F, Abbas W, Zia-Ur-Rehman M, Baig SA, Hashim M, Khan A & Rehman Hu (2021). Effect of green human resource management practices on organizational sustainability: the mediating role of environmental and employee performance. *Environmental Science and Pollution Research*, 28, 28191-28206.
- Amjad MS, Zahoor U, Bussmann RW, Altaf M, Gardazi SMH & Abbasi AM (2020). Ethnobotanical survey of the medicinal flora of Harighal, Azad Jammu & Kashmir, Pakistan. *Journal of ethnobiology and ethnomedicine*, 16, 1-28.
- Anderson VA, Anderson P, Northam E, Jacobs R & Catroppa C (2001). Development of executive functions through late childhood and adolescence in an Australian sample. *Developmental neuropsychology*, 20(1), 385-406.
- Andrés CMC, Pérez de la Lastra JM, Juan CA, Plou FJ & Pérez-Lebeña E (2024). Antioxidant Metabolism Pathways in Vitamins, Polyphenols, and Selenium: Parallels and Divergences. *International journal of molecular sciences*, 25(5), 2600.
- Anet A, Olakkaran S, Purayil AK & Puttaswamygowda GH (2019). Bisphenol A induced oxidative stress mediated genotoxicity in *Drosophila melanogaster*. *Journal of hazardous materials*, 370, 42-53.
- Anjum S, Rahman S, Kaur M, Ahmad F, Rashid H, Ansari RA & Raisuddin S (2011). Melatonin ameliorates bisphenol A-induced biochemical toxicity in testicular mitochondria of mouse. *Food and chemical toxicology*, 49(11), 2849-2854.
- Babu S, Uppu S, Claville MO & Uppu RM (2013). Prooxidant actions of bisphenol A (BPA) phenoxyl radicals: implications to BPA-related oxidative stress and toxicity. *Toxicology mechanisms and methods*, 23(4), 273-280.
- Bach MX, Minh TN, Anh DTN, Anh HN, Anh LV, Trung NQ, Xuan TD (2022). Protection and Rehabilitation Effects of Cordyceps militaris Fruit Body Extract and Possible Roles of Cordycepin and Adenosine. *Compounds*, 2(4), 388-403.
- Balogh GT, Illés J, Székely Z, Forrai E & Gere A (2003). Effect of different metal ions on the oxidative damage and antioxidant capacity of hyaluronic acid. *Archives of Biochemistry and Biophysics*, 410(1), 76-82.
- Bameri B, Armandeh M, Haghi-Aminjan H, Rahimifard M, Hassani S, Gholami M, Hooshangi Shayesteh MR (2024). Insights into the role of melatonin on the reproductive system toxicity induced by bisphenol A: a systematic review of preclinical studies. *Toxin reviews*, 43(1), 137-156.
- Bansal AK & Bilaspuri G (2011). Impacts of oxidative stress and antioxidants on semen functions. *Veterinary medicine international*, 2011(1), 686137.
- Barati E, Nikzad H & Karimian M (2020). Oxidative stress and male infertility: current knowledge of pathophysiology and role of antioxidant therapy in disease management. *Cellular and Molecular Life Sciences*, 77, 93-113.
- Bariar B Vestal CG & Richardson C (2013). Long-term effects of chromatin remodeling and DNA damage in stem cells induced by environmental and dietary agents. *Journal of Environmental Pathology, Toxicology and Oncology*, 32(4).
- Bayliak MM, Gospodaryov DV & Lushchak VI (2023). Homeostasis of carbohydrates and reactive oxygen species is critically changed in the brain of middle-aged mice: Molecular mechanisms and functional reasons. *BBA advances*, 3, 100077.
- Bilgi A, Abalı R, Bilgi PT, Şahin M, Tunçdemir M & Boran AB (2019). The apoptotic effects of bisphenol A exposure on the rat ovary: an experimental study. *Environmental Science and Pollution Research*, 26, 10198-10203.
- Bisht S, Faiq M, Tolahunase M & Dada R (2017). Oxidative stress and male infertility. *Nature Reviews Urology*, 14(8), 470-485.
- Bouayed J & Bohn T (2010). Exogenous antioxidants—Double-edged swords in cellular redox state: Health beneficial effects at physiologic doses versus deleterious effects at high doses. *Oxidative medicine and cellular longevity*, 3(4), 228-237.
- Bradford JB, Schlaepfer DR, Lauenroth WK & Palmquist KA (2020). Robust ecological drought projections for drylands in the 21st century. *Global change biology*, 26(7), 3906-3919.
- Brennan TH, Lewis LK, Gordon SJ & Prichard I (2024). Effectiveness of interventions to prevent or reverse pre-frailty and frailty in middle-aged community dwelling adults: A systematic review. *Preventive Medicine*, 108008.
- Burton GJ & Jauniaux E (2011). Oxidative stress. *Best practice & research Clinical obstetrics & gynaecology*, 25, 287-299.
- Cariati F, Carbone L, Conforti A, Bagnulo F, Peluso SR, Carotenuto C, Strina I (2020). Bisphenol A-induced epigenetic changes and its effects on the male reproductive system. *Frontiers in Endocrinology*, 11, 453.
- Carneiro CFD, da Costa GG, Neves K, Abreu MB, Tan PB, Rayêe D, Malički M (2023). Characterization of comments about bioRxiv and medRxiv preprints. *JAMA Network Open*, 6(8), e2331410-e2331410.
- Carneiro MFH & Colaiácovo MP (2023). Beneficial antioxidant effects of Coenzyme Q10 on reproduction. *Vitamins and Hormones*, 121, 143-167.
- Casals-Casas C & Desvergne B (2011). Endocrine disruptors: from endocrine to metabolic disruption. *Annual review of physiology*, 73(1), 135-162.
- Cetin I, Berti C & Calabrese S (2010). Role of micronutrients in the periconceptional period. *Human reproduction update*, 16(1), 80-95.
- Chattopadhyay S, Sahoo D, Subudhi U & Chainy G (2007). Differential expression profiles of antioxidant enzymes and glutathione redox status in hyperthyroid rats: a temporal analysis. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 146(3), 383-391.
- Chen FF (2007). Sensitivity of goodness of fit indexes to lack of measurement invariance. *Structural equation modeling: a multidisciplinary journal*, 14(3), 464-504.
- Chen S, Wang Z, Zhang M, Shi X, Wang L, An W, Yang L (2023). Chen et al. *Carbon Energy*, 5(8).
- Chianese R, Troisi J, Richards S, Scafuro M, Fasano S, Guida M, Meccariello R (2018). Bisphenol A in

- reproduction: epigenetic effects. *Current medicinal chemistry*, 25(6), 748-770.
- Choi YJ, Ha KH & Kim DJ (2017). Exposure to bisphenol A is directly associated with inflammation in healthy Korean adults. *Environmental Science and Pollution Research*, 24, 284-290.
- Cilio S, Rienzo M, Villano G, Mirto BF, Giampaglia G, Capone F, Crocetto F (2022). Beneficial effects of antioxidants in male infertility management: A narrative review. *Oxygen*, 2(1), 1-11.
- Cimmino I, Fiory F, Perruolo G, Miele C, Beguinot F, Formisano P & Oriente F. (2020). Potential mechanisms of bisphenol A (BPA) contributing to human disease. *International journal of molecular sciences*, 21(16), 5761.
- Cloughesy TF, Mochizuki AY, Orpilla JR, Hugo W, Lee AH, Davidson TB, Sanders CM (2019). Neoadjuvant anti-PD-1 immunotherapy promotes a survival benefit with intratumoral and systemic immune responses in recurrent glioblastoma. *Nature medicine*, 25(3), 477-486.
- Comminos AN, Jayasena CN & Dhillon WS (2014). The relationship between gut and adipose hormones, and reproduction. *Human reproduction update*, 20(2), 153-174.
- Costa, B., & Mercadal JG (1995). Costa: St. Martin's Press.
- Dattilo M, Giuseppe DA, Ettore C & Ménézo Y (2016). Improvement of gamete quality by stimulating and feeding the endogenous antioxidant system: mechanisms, clinical results, insights on gene-environment interactions and the role of diet. *Journal of assisted reproduction and genetics*, 33(12), 1633-1648.
- De Flora S, Micale RT, La Maestra S, Izzotti A, D'Agostini F, Camoirano A, Davalli P (2011). Upregulation of clusterin in prostate and DNA damage in spermatozoa from bisphenol A-treated rats and formation of DNA adducts in cultured human prostatic cells. *Toxicological Sciences*, 122(1), 45-51.
- de Haan G, Weersing E, Dontje B, van Os R, Bystrikh LV, Vellenga E & Miller G (2003). In vitro generation of long-term repopulating hematopoietic stem cells by fibroblast growth factor-1. *Developmental cell*, 4(2), 241-251.
- De Toni L, De Rocco Ponce M, Petre GC, Rtibi K, Di Nisio A & Foresta C (2020). Bisphenols and male reproductive health: from toxicological models to therapeutic hypotheses. *Frontiers in Endocrinology*, 11, 301.
- Dehghani MA, Sabahi A, Vanani AR, Amirrajab N, Ansari AC, Bazmi A & Alikarami K (2023). Mechanisms of bisphenol A toxicity in mitochondria. *Mitochondrial Intoxication* (pp. 409-423): Elsevier.
- Delbes G, Blázquez M, Fernandino JI, Grigorova P, Hales B, Metcalfe C, Rwigemera A (2022). Effects of endocrine disrupting chemicals on gonad development: Mechanistic insights from fish and mammals. *Environmental research*, 204, 112040.
- Dias MSdA, Parente JRF, Vasconcelos MIO & Dias FAC (2014). Intersetorialidade e Estratégia Saúde da Família: tudo ou quase nada a ver? *Ciência & Saúde Coletiva*, 19, 4371-4382.
- Dutta S, Majzoub A & Agarwal A (2019). Oxidative stress and sperm function: A systematic review on evaluation and management. *Arab journal of urology*, 17(2), 87-97.
- Edmonds DA, Caldwell RL, Brondizio ES & Siani SM (2020). Coastal flooding will disproportionately impact people on river deltas. *Nature communications*, 11(1), 4741.
- Ehrlich M & Lacey M (2013). DNA methylation and differentiation: silencing, upregulation and modulation of gene expression. *Epigenomics*, 5(5), 553-568.
- El Henafy HM, Ibrahim MA, Abd El Aziz SA & Gouda EM (2020). Oxidative Stress and DNA methylation in male rat pups provoked by the transplacental and translactational exposure to bisphenol A. *Environmental Science and Pollution Research*, 27, 4513-4519.
- Elswefy SES, Abdallah FR, Wahba AS, Hasan RA & Atteia HH (2020). Antifibrotic effect of curcumin, N-acetyl cysteine and propolis extract against bisphenol A-induced hepatotoxicity in rats: Prophylaxis versus co-treatment. *Life Sciences*, 260, 118245.
- Essawy AE, Jimmiey E M, Abdel-Wahab WM, Ali RG, Eweda SM & Abdou HM (2025). The protective efficacy of omega-3 polyunsaturated fatty acids on oxidative stress, inflammation, neurotransmitter perturbations, and apoptosis induced by monosodium glutamate in the brain of male rats. *Metabolic Brain Disease*, 40(1), 1-14.
- Evans A (2022). Characterizing the relative biologic effectiveness following clinical and space-relevant ionizing radiation exposures and exploring curcumin nanoparticles as effective radiological countermeasures: Lawrence Livermore National Lab.(LLNL), Livermore, CA (United States).
- Feng S, Sun H, Yan X, Zhu H, Zou Z, Shen S & Liu HX (2023). Dense reinforcement learning for safety validation of autonomous vehicles. *Nature*, 615(7953), 620-627.
- Ferguson KK, Cantonwine DE, McElrath TF, Mukherjee B & Meeker JD (2016). Repeated measures analysis of associations between urinary bisphenol-A concentrations and biomarkers of inflammation and oxidative stress in pregnancy. *Reproductive toxicology*, 66, 93-98.
- Fialkow L, Wang Y & Downey GP (2007). Reactive oxygen and nitrogen species as signaling molecules regulating neutrophil function. *Free Radical Biology and Medicine*, 42(2), 153-164.
- Fuhrman VF, Tal A & Arnon S (2015). Why endocrine disrupting chemicals (EDCs) challenge traditional risk assessment and how to respond. *Journal of hazardous materials*, 286, 589-611.
- Funk IK & March FA (1897). *A standard dictionary of the English language* (Vol. 2): Funk & Wagnalls Company.
- Ganesan S & Keating AF (2016). Bisphenol A-induced ovotoxicity involves DNA damage induction to which the ovary mounts a protective response indicated by increased expression of proteins involved in DNA repair and xenobiotic biotransformation. *Toxicological Sciences*, 152, 169-180.

- Gassman NR, Coskun E, Stefanick DF, Horton JK, Jaruga P, Dizdaroglu M & Wilson SH (2015). Bisphenol A promotes cell survival following oxidative DNA damage in mouse fibroblasts. *PLoS One*, 10(2), e0118819.
- Geesink J & Meijer DK (2017). Cancer Is Promoted by Cellular States of Electromagnetic Decoherence and Can Be Corrected by Exposure to Coherent Non-Ionizing Electromagnetic Fields. A Physical Model about Cell-Sustaining and Cell-Decaying Soliton Eigen-Frequencies. A Phys. Model about cell-sustaining cell-decaying Solit. eigen-frequencies, 1-45.
- Gezer A, Üstündağ H, Kılıç Baygutalp N, Erbaş E & Özkaraca M (2024). The Protective Effect of Gallic Acid Against Bisphenol A-Induced Ovarian Toxicity and Endocrine Disruption in Female Rats. *Journal of medicinal food*, 27(7), 651-660.
- Ghimire S, Shrestha A, Yadav UN, Mistry SK, Chapadia B, Yadav OP, Mehata S (2022). Older adults with pre-existing noncommunicable conditions and their healthcare access amid COVID-19 pandemic: a cross-sectional study in eastern Nepal. *BMJ open*, 12(2), e056342.
- Gignac GE & Zajenkowski M (2023). Still no Dunning-Kruger effect: A reply to Hiller. *Intelligence*, 97, 101733.
- Graille M, Wild P, Sauvain JJ, Hemmendinger M, Guseva Canu I & Hopf NB (2020). Urinary 8-OHdG as a biomarker for oxidative stress: a systematic literature review and meta-analysis. *International journal of molecular sciences*, 21(11), 3743.
- Hahladakis JN, Iacovidou E & Gerassimidou S (2023). An overview of the occurrence, fate, and human risks of the bisphenol-A present in plastic materials, components, and products. *Integrated environmental assessment and management*, 19(1), 45-62.
- Hajimehdipoor H, Kondori BM, Amin GR, Adib N, Rastegar H & Shekarchi M (2012). Development of a validated HPLC method for the simultaneous determination of flavonoids in *Cuscuta chinensis* Lam. by ultra-violet detection. *DARU Journal of Pharmaceutical Sciences*, 20, 1-6.
- Heidarzadi S, Rashidi Z, Jalili C & Gholami M (2022). Overview of biological effects of melatonin on testis: A review. *Andrologia*, 54(11), e14597.
- Heindel JJ, Blumberg B, Cave M, Machtinger R, Mantovani A, Mendez MA, Sargis R (2017). Metabolism disrupting chemicals and metabolic disorders. *Reproductive toxicology*, 68, 3-33.
- Hengstler J, Foth H, Gebel T, Kramer PJ, Lilienblum W, Schweinfurth H, Gundert-Remy U (2011). Critical evaluation of key evidence on the human health hazards of exposure to bisphenol A. *Critical reviews in toxicology*, 41(4), 263-291.
- Hernández-Silva CD, Villegas-Pineda JC & Pereira-Suárez AL (2020). Expression and role of the G protein-coupled estrogen receptor (GPR30/GPER) in the development and immune response in female reproductive cancers. *Frontiers in Endocrinology*, 11, 544.
- Herrero JL, Roberts M, Delicato LS, Gieselmann MA, Dayan P & Thiele A (2008). Acetylcholine contributes through muscarinic receptors to attentional modulation in V1. *Nature*, 454(7208), 1110-1114.
- Huang Y, Yu L, Lu P, Wei Y, Fu L, Hou J, Chen L (2022). Evaluate the bisphenol A-induced redox state in cells, zebrafish and in vivo with a hydrogen peroxide turn-on fluorescent probe. *Journal of hazardous materials*, 424, 127425.
- Huc L, Lemarié A, Guéraud F & Héliers-Toussaint C (2012). Low concentrations of bisphenol A induce lipid accumulation mediated by the production of reactive oxygen species in the mitochondria of HepG2 cells. *Toxicology in Vitro*, 26(5), 709-717.
- Hwang B, Lee JH & Bang D (2018). Single-cell RNA sequencing technologies and bioinformatics pipelines. *Experimental & molecular medicine*, 50(8), 1-14.
- Jain SK, Stevens CM, Margret JJ & Levine SN (2024). Alzheimer's Disease: A Review of Pathology, Current Treatments, and the Potential Therapeutic Effect of Decreasing Oxidative Stress by Combined Vitamin D and L-Cysteine Supplementation. *Antioxidants & redox signaling*, 40(10-12), 663-678.
- Jędrejko KJ, Lazur J & Muszyńska B (2021). *Cordyceps militaris*: An overview of its chemical constituents in relation to biological activity. *Foods*, 10(11), 2634.
- Jeremias G, Gonçalves FJ, Pereira JL & Asselman J (2020). Prospects for incorporation of epigenetic biomarkers in human health and environmental risk assessment of chemicals. *Biological Reviews*, 95(3), 822-846.
- Justin Margret J & Jain SK (2024). The association between alpha-1 antitrypsin and B-type natriuretic peptide blood levels in healthy African Americans. *BMC Research Notes*, 17(1), 331.
- Kamp TJ & Hell JW (2000). Regulation of cardiac L-type calcium channels by protein kinase A and protein kinase C. *Circulation research*, 87(12), 1095-1102.
- Kang J, Liu L, Liu M, Wu X & Li J (2018). Antibacterial activity of gallic acid against *Shigella flexneri* and its effect on biofilm formation by repressing mdoH gene expression. *Food Control*, 94, 147-154.
- Kaur SP & Gupta V (2020). COVID-19 Vaccine: A comprehensive status report. *Virus research*, 288, 198114.
- Keane KN, Cruzat VF, Carlessi R, de Bittencourt Jr PIH & Newsholme P (2015). Molecular events linking oxidative stress and inflammation to insulin resistance and β -cell dysfunction. *Oxidative medicine and cellular longevity*, 2015(1), 181643.
- Khalaf A, Ahmed W, Moselhy W, Abdel-Halim B & Ibrahim M (2019). Protective effects of selenium and nano-selenium on bisphenol-induced reproductive toxicity in male rats. *Human & experimental toxicology*, 38(4), 398-408.
- Khan S, Beigh S, Chaudhari BP, Sharma S, Aliul Hasan Abdi S, Ahmad S, Raisuddin S (2016). Mitochondrial dysfunction induced by Bisphenol A is a factor of its hepatotoxicity in rats. *Environmental Toxicology*, 31(12), 1922-1934.
- Kini RD, Kumar S, Nayanatara A & Megha G (2024). A study on bisphenol induce renal damage and role of vitamin c as an antioxidant on bisphenol induced renal damage in wistar rats. *La Clinica Terapeutica*, 175(6).

- Kopustinskiene DM & Bernatoniene J (2021). Molecular mechanisms of melatonin-mediated cell protection and signaling in health and disease. *Pharmaceutics*, 13(2), 129.
- Krylatov AV, Maslov LN, Voronkov NS, Boshchenko AA, Popov SV, Gomez L, Downey JM (2018). Reactive oxygen species as intracellular signaling molecules in the cardiovascular system. *Current cardiology reviews*, 14(4), 290-300.
- Kumar S & Pandey A (2015). Free radicals: health implications and their mitigation by herbals. *British Journal of Medicine and Medical Research*, 7(6), 438-457.
- Kunysz M, Mora-Janiszewska O & Darmochwał-Kolarz D (2021). Epigenetic modifications associated with exposure to endocrine disrupting chemicals in patients with gestational diabetes mellitus. *International journal of molecular sciences*, 22(9), 4693.
- La Vignera S & Vita R (2018). Thyroid dysfunction and semen quality. *International Journal of Immunopathology and Pharmacology*, 32, 2058738418775241.
- Langie SA, Koppen G, Desaulniers D, Al-Mulla F, Al-Temaimi R, Amedei A, Brunborg G (2015). Causes of genome instability: the effect of low dose chemical exposures in modern society. *Carcinogenesis*, 36(Suppl_1), S61-S88.
- Li Y, Dong Z, Liu S, Gao F, Zhang J, Peng Z, Pan X (2022). Astaxanthin improves the development of the follicles and oocytes through alleviating oxidative stress induced by BPA in cultured follicles. *Scientific Reports*, 12(1), 7853.
- Lind S, Ingvaldsen RB & Furevik T (2018). Arctic warming hotspot in the northern Barents Sea linked to declining sea-ice import. *Nature climate change*, 8(7), 634-639.
- Liu P, Yuan W, Fu J, Jiang Z, Hayashi H & Neubig G (2023). Pre-train, prompt, and predict: A systematic survey of prompting methods in natural language processing. *ACM Computing Surveys*, 55(9), 1-35.
- Liu X, Miao M, Zhou Z, Gao E, Chen J, Wang J, Li DK (2015). Exposure to bisphenol-A and reproductive hormones among male adults. *Environmental toxicology and pharmacology*, 39(2), 934-941.
- Lombó M, Fernández-Díez C, González-Rojo S & Herráez MP (2019). Genetic and epigenetic alterations induced by bisphenol A exposure during different periods of spermatogenesis: from spermatozoa to the progeny. *Scientific Reports*, 9(1), 18029.
- Loren P, Sánchez R, Arias ME, Felmer R, Risopatrón J & Cheuquemán C (2017). Melatonin scavenger properties against oxidative and nitrosative stress: impact on gamete handling and in vitro embryo production in humans and other mammals. *International journal of molecular sciences*, 18(6), 1119.
- Lu J, Wang Y, Li J, Mao L, Nguyen SH, Duarte T, Guo J (2018). Triclosan at environmentally relevant concentrations promotes horizontal transfer of multidrug resistance genes within and across bacterial genera. *Environment international*, 121, 1217-1226.
- Mączka W, Grabarczyk M & Wińska K (2022). Can antioxidants reduce the toxicity of bisphenol? *Antioxidants*, 11(2), 413.
- Malivindi R, Rago V, De Rose D, Gervasi MC, Cione E, Russo G, Aquila S (2018). Influence of all-trans retinoic acid on sperm metabolism and oxidative stress: Its involvement in the physiopathology of varicocele-associated male infertility. *Journal of Cellular Physiology*, 233(12), 9526-9537.
- Marcu D, Keyser S, Petrik L, Fuhrmann S & Maree L (2023). Contaminants of emerging concern (CECs) and male reproductive health: challenging the future with a double-edged sword. *Toxics*, 11(4), 330.
- Marí M, Morales A, Colell A, García-Ruiz C & Fernández-Checa JC (2009). Mitochondrial glutathione, a key survival antioxidant. *Antioxidants & redox signaling*, 11(11), 2685-2700.
- Martos YM, Catalán M, Jordan TA, Golynsky A, Golynsky D, Eagles G & Vaughan DG (2017). Heat flux distribution of Antarctica unveiled. *Geophysical Research Letters*, 44(22), 11,417-411,426.
- Matarrese P, Mattia G, Pagano M T, Pontecorvi G, Ortona E, Malorni W & Carè A (2021). The sex-related interplay between time and cancer: On the critical role of estrogen, micromRNAs and autophagy. *Cancers*, 13(13), 3287.
- Mathur S, Metcalfe T, Woitaszek M, Bruntt H, Verner G, Christensen-Dalsgaard J, Karoff C (2012). A uniform asteroseismic analysis of 22 solar-type stars observed by Kepler. *The Astrophysical Journal*, 749(2), 152.
- Matuszczak E, Komarowska MD, Debek W & Hermanowicz A (2019). The impact of bisphenol A on fertility, reproductive system, and development: a review of the literature. *International journal of endocrinology*, 2019(1), 4068717.
- McKay NS, Gordon BA, Hornbeck RC, Dincer A, Flores S, Keefe SJ, Millar PR (2023). Positron emission tomography and magnetic resonance imaging methods and datasets within the Dominantly Inherited Alzheimer Network (DIAN). *Nature neuroscience*, 26(8), 1449-1460.
- Megha K, Arathi A, Shikha S, Alka R, Ramya P & Mohanan P (2024). Significance of melatonin in the regulation of circadian rhythms and disease management. *Molecular Neurobiology*, 1-31.
- Meli R, Monnolo A, Annunziata C, Pirozzi C & Ferrante MC (2020). Oxidative stress and BPA toxicity: an antioxidant approach for male and female reproductive dysfunction. *Antioxidants*, 9(5), 405.
- Mendiola J, Jørgensen N, Andersson AM, Calafat AM, Ye X, Redmon JB, Thurston SW (2010). Are environmental levels of bisphenol A associated with reproductive function in fertile men? *Environmental health perspectives*, 118(9), 1286-1291.
- Micheli L, Cerretani D, Collodel G, Menchiari A, Moltoni L, Fiaschi A & Moretti E (2016). Evaluation of enzymatic and non-enzymatic antioxidants in seminal plasma of men with genitourinary infections, varicocele and idiopathic infertility. *Andrology*, 4(3), 456-464.
- Mileva G, Baker SL, Konkle AT & Bielajew C (2014). Bisphenol-A: epigenetic reprogramming and effects on reproduction and behavior. *International Journal*

- of Environmental Research and Public Health, 11(7), 7537-7561.
- Mocchegiani E, Bulian D, Santarelli L, Tibaldi A, Muzzioli M, Pierpaoli W & Nicola F (1994). The immuno-reconstituting effect of melatonin or pineal grafting and its relation to zinc pool in aging mice. *Journal of neuroimmunology*, 53(2), 189-201.
- Mohammed ET, Hashem KS, Ahmed AE, Aly MT, Aleya L & Abdel-Daim MM (2020). Ginger extract ameliorates bisphenol A (BPA)-induced disruption in thyroid hormones synthesis and metabolism: involvement of Nrf-2/HO-1 pathway. *Science of the total environment*, 703, 134664.
- Molina-López AM, Bujalance-Reyes F, Ayala-Soldado N, Mora-Medina R, Lora-Benítez A & Moyano-Salvago R (2023). An overview of the health effects of bisphenol a from a one health perspective. *Animals*, 13(15), 2439.
- Mondal S & Bandyopadhyay A (2024). Antioxidants in mitigating phthalate-induced male reproductive toxicity: A comprehensive review. *Chemosphere*, 143297.
- Mou D, Wang J, Liu H, Chen Y, Che L, Fang Z, Li J (2018). Maternal methyl donor supplementation during gestation counteracts bisphenol A-induced oxidative stress in sows and offspring. *Nutrition*, 45, 76-84.
- Nadal A, Quesada I, Tuduri E, Nogueiras R & Alonso-Magdalena P (2017). Endocrine-disrupting chemicals and the regulation of energy balance. *Nature Reviews Endocrinology*, 13(9), 536-546.
- Nagarajan M, Maadurshni GB & Manivannan J (2024). Bisphenol a (BPA) exposure aggravates hepatic oxidative stress and inflammatory response under hypertensive milieu—impact of low dose on hepatocytes and influence of MAPK and ER stress pathways. *Food and chemical toxicology*, 183, 114197.
- Navarro-Martín L, Martyniuk CJ & Mennigen JA (2020). Comparative epigenetics in animal physiology: an emerging frontier. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 36, 100745.
- Nelson CA, Bhutta ZA, Harris NB, Danese A & Samara M (2020). Adversity in childhood is linked to mental and physical health throughout life. *bmj*, 371.
- Nordberg A, Rinne JO, Kadir A & Långström B (2010). The use of PET in Alzheimer disease. *Nature Reviews Neurology*, 6(2), 78-87.
- Ola-Davies OE & Olukole SG (2018). Gallic acid protects against bisphenol A-induced alterations in the cardio-renal system of Wistar rats through the antioxidant defense mechanism. *Biomedicine & Pharmacotherapy*, 107, 1786-1794.
- Olukole SG, Lanipekun DO, Ola-Davies EO & Oke BO (2019). Maternal exposure to environmentally relevant doses of bisphenol A causes reproductive dysfunction in F1 adult male rats: protective role of melatonin. *Environmental Science and Pollution Research*, 26, 28940-28950.
- Onuzulu CD, Rotimi OA & Rotimi SO (2019). Epigenetic modifications associated with in utero exposure to endocrine disrupting chemicals BPA, DDT and Pb. *Reviews on environmental health*, 34(4), 309-325.
- Ornoy A, Reece EA, Pavlinkova G, Kappen C & Miller RK (2015). Effect of maternal diabetes on the embryo, fetus, and children: congenital anomalies, genetic and epigenetic changes and developmental outcomes. *Birth Defects Research Part C: Embryo Today: Reviews*, 105(1), 53-72.
- Ourique GM, Saccol EM, Pês TS, Glanzner WG, Schiefelbein SH, Woehl VM, Barreto KP (2016). Protective effect of vitamin E on sperm motility and oxidative stress in valproic acid treated rats. *Food and chemical toxicology*, 95, 159-167.
- Paulose T, Speroni L, Sonnenschein C & Soto AM (2015). Estrogens in the wrong place at the wrong time: Fetal BPA exposure and mammary cancer. *Reproductive toxicology*, 54, 58-65.
- Perera L, Li Y, Coons LA, Houtman R, van Beuningen R, Goodwin B, Teng CT (2017). Binding of bisphenol A, bisphenol AF, and bisphenol S on the androgen receptor: Coregulator recruitment and stimulation of potential interaction sites. *Toxicology in Vitro*, 44, 287-302.
- Peretz J, Vrooman L, Ricke WA, Hunt PA, Ehrlich S, Hauser R, VandeVoort CA (2014). Bisphenol A and reproductive health: update of experimental and human evidence, 2007–2013. *Environmental health perspectives*, 122(8), 775-786.
- Peris E, Micallef P, Paul A, Palsdottir V, Enejder A, Bauzá-Thorbrügge M, Asterholm IW (2019). Antioxidant treatment induces reductive stress associated with mitochondrial dysfunction in adipocytes. *Journal of Biological Chemistry*, 294(7), 2340-2352.
- Phillips DH & Arlt VM (2009). Genotoxicity: damage to DNA and its consequences. *Molecular, Clinical and Environmental Toxicology: Volume 1: Molecular Toxicology*, 87-110.
- Pirozzi C, Lama A, Annunziata C, Cavaliere G, Ruiz-Fernandez C, Monnolo A, Mollica MP (2020). Oral bisphenol A worsens liver immune-metabolic and mitochondrial dysfunction induced by high-fat diet in adult mice: cross-talk between oxidative stress and inflammasome pathway. *Antioxidants*, 9(12), 1201.
- Pivonello C, Muscogiuri G, Nardone A, Garifalos F, Provisiero DP, Verde N, Auriemma RS (2020). Bisphenol A: an emerging threat to female fertility. *Reproductive Biology and Endocrinology*, 18, 1-33.
- Quan C, Wang C, Duan P, Huang W & Yang K (2017). Prenatal bisphenol a exposure leads to reproductive hazards on male offspring via the Akt/mTOR and mitochondrial apoptosis pathways. *Environmental Toxicology*, 32(3), 1007-1023.
- Rakha SI, Elmetwally MA, El-Sheikh Ali H, Balboula A, Mahmoud AM & Zaabel SM (2022). Importance of antioxidant supplementation during in vitro maturation of mammalian oocytes. *Veterinary Sciences*, 9(8), 439.
- Ramaswamy S & Weinbauer GF (2014). Endocrine control of spermatogenesis: Role of FSH and LH/testosterone. *Spermatogenesis*, 4(2), e996025.
- Ramos E, Lopez-Munoz F, Gil-Martin E, Egea J, Alvarez-Merz I, Painuli S, Romero A (2021). The coronavirus disease 2019 (COVID-19): key emphasis on melatonin safety and therapeutic efficacy. *Antioxidants*, 10(7), 1152.

- Rejhová A, Opattová A, Čumová A, Slíva D & Vodička P (2018). Natural compounds and combination therapy in colorectal cancer treatment. *European journal of medicinal chemistry*, 144, 582-594.
- Ribeiro LS, de Melo Orfao JJ & Pereira MFR (2017). Direct catalytic production of sorbitol from waste cellulosic materials. *Bioresource technology*, 232, 152-158.
- Rizzo R, Bortolotti D, Rizzo S & Schiuma G (2023). Endocrine Disruptors, Epigenetic Changes, and Transgenerational Transmission Environment Impact on Reproductive Health: A Translational Approach (pp. 49-74): Springer International Publishing Cham.
- Rosa J & Flores N (2017). Unsettling race and language: Toward a raciolinguistic perspective. *Language in society*, 46(5), 621-647.
- Rosenfeld CS, Sieli PT, Warzak DA, Ellersieck MR, Pennington KA & Roberts RM (2013). Maternal exposure to bisphenol A and genistein has minimal effect on Avy/a offspring coat color but favors birth of agouti over nonagouti mice. *Proceedings of the National Academy of Sciences*, 110(2), 537-542.
- Ryu DY, Pang WK, Adegoke EO, Rahman MS, Park YJ & Pang MG (2023). Bisphenol-A disturbs hormonal levels and testis mitochondrial activity, reducing male fertility. *Human Reproduction Open*, 2023(4), hoad044.
- Sadek KM, Khalifa NE, Alshial EE, Abdelnour SA, Mohamed AAR & Noreldin AE (2024). Potential hazards of bisphenol A on the male reproductive system: Induction of programmed cell death in testicular cells. *Journal of Biochemical and Molecular Toxicology*, 38(9), e23844.
- Sahu A & Verma R (2023). Bisphenol S dysregulates thyroid hormone homeostasis; Testicular survival, redox and metabolic status: Ameliorative actions of melatonin. *Environmental toxicology and pharmacology*, 104, 104300.
- Salim A, Pangkahila WI & Sriwidayani NP (2023). The Administration of Cordyceps Militaris Inhibited the Decrease of Leydig Cell Number and Testosterone Level in Male Wistar Rats (*Rattus Norvegicus*) Exposed with Cigarette Smoke. *European Journal of Biomedical Research*, 2(2), 1-4.
- Salvador R, Barros MV, Da Luz LM, Piekarski CM & de Francisco AC (2020). Circular business models: Current aspects that influence implementation and unaddressed subjects. *Journal of Cleaner Production*, 250, 119555.
- Sánchez-Cruzado C, Santiago Campión R & Sánchez-Compañía MT (2021). Teacher digital literacy: The indisputable challenge after COVID-19. *Sustainability*, 13(4), 1858.
- Santiago I, Moreno-Munoz A, Quintero-Jiménez P, Garcia-Torres F & Gonzalez-Redondo MJ (2021). Electricity demand during pandemic times: The case of the COVID-19 in Spain. *Energy policy*, 148, 111964.
- Sarg NH, Zaher DM, Jayab NNA, Mostafa SH, Ismail HH & Omar HA (2024). The interplay of p38 MAPK signaling and mitochondrial metabolism, a dynamic target in cancer and pathological contexts. *Biochemical Pharmacology*, 116307.
- Sedha S, Kumar S & Shukla S (2015). Role of oxidative stress in male reproductive dysfunctions with reference to phthalate compounds. *Urology journal*, 12(5), 2304-2316.
- Shah MS & Brownlee M (2016). Molecular and cellular mechanisms of cardiovascular disorders in diabetes. *Circulation research*, 118(11), 1808-1829.
- Sharma M, Sharma R, Gupta P & Srivastava S (2019). Bisphenol-A induced oxidative stress and its fertility aspects. *International Journal of Pharmaceutical Sciences and Research*, 10, 3519-3531.
- Sifakis S, Androutopoulos VP, Tsatsakis AM & Spandidos DA (2017). Human exposure to endocrine disrupting chemicals: effects on the male and female reproductive systems. *Environmental toxicology and pharmacology*, 51, 56-70.
- Singh CK, Chhabra G, Ndiaye MA, Garcia-Peterson LM, Mack NJ & Ahmad N (2018). The role of sirtuins in antioxidant and redox signaling. *Antioxidants & redox signaling*, 28(8), 643-661.
- Skakkebaek NE, Rajpert-De Meyts E, Buck Louis GM, Toppari J, Andersson AM, Eisenberg ML, Sapra KJ (2016). Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiological reviews*, 96(1), 55-97.
- Spahis S, Borys JM & Levy E (2017). Metabolic syndrome as a multifaceted risk factor for oxidative stress. *Antioxidants & redox signaling*, 26(9), 445-461.
- Srivastava N, Mansimov E & Salakhudinov R (2015). Unsupervised learning of video representations using lstms. Paper presented at the International conference on machine learning.
- Stavridis K, Triantafyllidou O, Pisimisi M & Vlahos N (2022). Bisphenol-A and female fertility: an update of existing epidemiological studies. *Journal of Clinical Medicine*, 11(23), 7227.
- Steffensen V (2020). Fire country: How Indigenous fire management could help save Australia: CSIRO Publishing.
- Sun Q, Li Y, Shi L, Hussain R, Mehmood K, Tang Z & Zhang H (2022). Heavy metals induced mitochondrial dysfunction in animals: Molecular mechanism of toxicity. *Toxicology*, 469, 153136.
- Tain YL & Hsu CN (2024). Melatonin Use during Pregnancy and Lactation Complicated by Oxidative Stress: Focus on Offspring's Cardiovascular-Kidney-Metabolic Health in Animal Models. *Antioxidants*, 13(2), 226.
- Tan DX, Manchester LC, Terron MP, Flores LJ & Reiter RJ (2007). One molecule, many derivatives: a never-ending interaction of melatonin with reactive oxygen and nitrogen species? *Journal of pineal research*, 42(1), 28-42.
- Tiberi J, Cesarini V, Stefanelli R, Canterini S, Fiorenza MT & La Rosa P (2023). Sex differences in antioxidant defence and the regulation of redox homeostasis in physiology and pathology. *Mechanisms of ageing and development*, 211, 111802.
- Tiwari B, Sellamuthu B, Ouarda Y, Drogui P, Tyagi RD & Buelna G (2017). Review on fate and mechanism of removal of pharmaceutical pollutants from

- wastewater using biological approach. *Bioresource technology*, 224, 1-12.
- Toporova L & Balaguer P (2020). Nuclear receptors are the major targets of endocrine disrupting chemicals. *Molecular and cellular endocrinology*, 502, 110665.
- Udeochu JC, Amin S, Huang Y, Fan L, Torres ERS, Carling GK, Kauwe G (2023). Tau activation of microglial cGAS-IFN reduces MEF2C-mediated cognitive resilience. *Nature neuroscience*, 26(5), 737-750.
- Ursini F, Maiorino M & Forman HJ (2016). Redox homeostasis: The Golden Mean of healthy living. *Redox biology*, 8, 205-215.
- Valko M Rhodes C Moncol J, Izakovic M & Mazur M (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-biological interactions*, 160(1), 1-40.
- Vallese A, Cordone V, Pecorelli A & Valacchi G (2023). Ox-inflammasome involvement in neuroinflammation. *Free Radical Biology and Medicine*.
- Varshavsky J, Smith A, Wang A, Hom E, Izano M, Huang H, Woodruff TJ (2020). Heightened susceptibility: A review of how pregnancy and chemical exposures influence maternal health. *Reproductive toxicology*, 92, 14-56.
- Wang Y, Yin W & Zeng J (2019). Global convergence of ADMM in nonconvex nonsmooth optimization. *Journal of Scientific Computing*, 78, 29-63.
- Wetherill YB, Akingbemi BT, Kanno J, McLachlan JA, Nadal A, Sonnenschein C, Belcher SM (2007). In vitro molecular mechanisms of bisphenol A action. *Reproductive toxicology*, 24(2), 178-198.
- Wong CY, Boon-Itt S & Wong CW (2011). The contingency effects of environmental uncertainty on the relationship between supply chain integration and operational performance. *Journal of Operations management*, 29(6), 604-615.
- Xiao K, Zhai J, Feng Y, Zhou N, Zhang X, Zou JJ, Shen X (2020). Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins. *Nature*, 583(7815), 286-289.
- Yahav D, Franceschini E, Koppel F, Turjeman A, Babich T, Bitterman R, Eliakim-Raz N (2019) Seven versus 14 days of antibiotic therapy for uncomplicated gram-negative bacteremia: a noninferiority randomized controlled trial. *Clinical Infectious Diseases*, 69(7), 1091-1098.
- Yan YQ, Liu M, Xu ZJ, Xu ZJ, Huang YX, Li XM, Lei XG (2024). Optimum doses and forms of selenium maintaining reproductive health via regulating homeostasis of gut microbiota and testicular redox, inflammation, cell proliferation, and apoptosis in roosters. *The Journal of Nutrition*, 154(2), 369-380.
- Yin L, Dai Y, Cui Z, Jiang X, Liu W, Han F, Liu J (2017). The regulation of cellular apoptosis by the ROS-triggered PERK/EIF2 α /chop pathway plays a vital role in bisphenol A-induced male reproductive toxicity. *Toxicology and Applied Pharmacology*, 314, 98-108.
- Yousef M, Abdallah G & Kamel K (2003). Effect of ascorbic acid and vitamin E supplementation on semen quality and biochemical parameters of male rabbits. *Animal reproduction science*, 76, 99-111.
- Zahran E, Elbahnaswy S, Mamdouh AZ & El-Matbouli M (2021). Xenosteroids in aquaculture with special consideration to Lake Manzala (Northern delta lake, Egypt): Types, sources and mechanism of action. *Aquaculture Research*, 52(12), 5962-5977.
- Zhang M, Dai X, Lu Y, Miao Y, Zhou C, Cui Z, Xiong B (2017). Melatonin protects oocyte quality from Bisphenol A-induced deterioration in the mouse. *Journal of pineal research*, 62(3), e12396.
- Zhang, Y, Du Y, Le W, Wang K, Kieffer N & Zhang J (2011). Redox control of the survival of healthy and diseased cells. *Antioxidants & redox signaling*, 15(11), 2867-2908.
- Zhao Q, Liu Y, Wang X, Zhu Y, Jiao Y, Bao Y & Shi W (2023). *Cuscuta chinensis* flavonoids reducing oxidative stress of the improve sperm damage in bisphenol A exposed mice offspring. *Ecotoxicology and environmental safety*, 255, 114831.
- Zhou Z, Liu X, Sun K, Lin C, Ma J, He M & Ouyang W (2019). Persulfate-based advanced oxidation processes (AOPs) for organic-contaminated soil remediation: A review. *Chemical Engineering Journal*, 372, 836-851.
- Zimmet P, Boyko E, Collier G & de Courten M (1999). Etiology of the metabolic syndrome: potential role of insulin resistance, leptin resistance, and other players. *Annals of the New York Academy of Sciences*, 892(1), 25-44.